

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 1 364 657 A1**

(12)

**EUROPEAN PATENT APPLICATION**

published in accordance with Art. 158(3) EPC

(43) Date of publication:

**26.11.2003 Bulletin 2003/48**

(21) Application number: **02711350.5**

(22) Date of filing: **06.02.2002**

(51) Int Cl.7: **A61K 38/21, A61K 38/17, A61K 39/395, A61K 45/00, A61P 35/00, A61P 35/02, A61P 19/00, C12Q 1/02, C12Q 1/66, C12Q 1/68, C12N 15/12, C07K 14/705, C07K 16/28**

(86) International application number:

**PCT/JP02/00989**

(87) International publication number:

**WO 02/064159 (22.08.2002 Gazette 2002/34)**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR**

Designated Extension States:

**AL LT LV MK RO SI**

(30) Priority: **07.02.2001 JP 2001031492**

(83) Declaration under Rule 28(4) EPC (expert solution)

(71) Applicant: **CHUGAI SEIYAKU KABUSHIKI KAISHA**  
**Tokyo, 115-8543 (JP)**

(72) Inventors:

- **KOSAKA, Masaaki**  
**Tokushima-shi, Tokushima 770-8075 (JP)**
- **OZAKI, Shuji**  
**Tokushima-shi, Tokushima 770-0045 (JP)**
- **WAKAHARA, Yuji, C/O CHUGAI SEIYAKU K.K.**  
**Gotenba-shi, Shizuoka 412-8513 (JP)**

(74) Representative: **Harding, Charles Thomas et al**  
**D. Young & Co.**  
**21 New Fetter Lane**  
**London EC4A 1DA (GB)**

**(54) REMEDIES FOR TUMOR IN HEMATOPOIETIC ORGANS**

(57) An inducing agent or enhancing agent, for the expression of HM1.24 antigen in hematopoietic tumor cells, comprising interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein as an active ingredient, as well as an anti-tumor

agent for hematopoietic tumors which comprises a combination of said inducing agent or enhancing agent and an antibody against HM1.24.

**EP 1 364 657 A1**

**Description****FIELD OF THE INVENTION**

5 [0001] The present invention relates to the use of interferon  $\alpha$ , interferon  $\gamma$ , and the IRF-2 protein as expression-enhancing agents for HM1.24 antigen in myeloma.

[0002] The present invention also relates to the use of interferon  $\alpha$ , interferon  $\gamma$ , and the IRF-2 protein as expression-enhancing agents for HM1.24 antigen in lymphatic tumors. Furthermore, the present invention relates to a method of treating leukemia using anti-HM1.24 antibody and interferon  $\alpha$  or interferon  $\gamma$ .

**BACKGROUND ART**

15 [0003] Leukocytes occurring in normal human peripheral blood comprise granulocytes, monocytes and lymphocytes, and granulocytes are further divided into neutrophils, eosinophils, and basophils. In the production of these blood cells, myelocytic stem cells and lymphatic stem cells differentiate from common undifferentiated hematopoietic stem cells, and from these stem cells, finally, each line of leukocytes differentiates. Blood cell-related cells including these hematopoietic stem cells are also referred to as hematopoietic cells. Tumors (hematopoietic tumors) of hematopoietic cells include leukemia, lymphoma, myeloma, and the like.

20 [0004] Leukemia is a disease of cancerated hematopoietic cells, in which bone marrow is occupied by leukemic cells and thereby normal hematopoietic functions are inhibited, resulting in the decreased production of normal blood cells and in subsequent development of anemia, leukopenia and thrombocytopenia. Also, based on the origin of leukemic cells, leukemia is roughly divided into two types: myelocytic leukemia and lymphocytic leukemia, each of which is further divided into the acute form and the chronic form. Furthermore, as a subtype, mixed lineage leukemia having cellular traits of the two lineages, bone marrow lineage and lymphocyte lineage, is also known.

25 [0005] Tumorigenesis takes place at the level of hematopoietic stem cells, wherein there are a case in which differentiation stops at a certain stage of differentiation and maturation and tumors are only formed in the cells upstream thereof, and a case in which the functions of differentiation and maturation are retained though it evades the biological regulatory functions and exhibits autonomous growth. The former includes acute leukemia, and the latter includes chronic leukemia and the myelodysplastic syndrome. Based on the identity of the growing cells, acute leukemia is roughly divided into acute myelocytic leukemia (AML), acute monocytic leukemia (AMoL), acute erythroleukemia, megakaryoblastic leukemia, and acute lymphocytic leukemia (ALL).

30 [0006] As a subtype, acute promyelocytic leukemia (APL) is known. Acute leukemia and the myelodysplastic syndrome may be classified based on the French-American-British classification (FAB classification). In the FAB classification, acute lymphocytic leukemia is divided into L1, L2, and L3, and for example Burkitt's lymphoma is classified into L3. Acute myelocytic leukemia is divided into M0, M1, M2, M3, M4, M5, M6, and M7 and, for example, erythrocyte abnormality is classified into M6 and megakaryoblastic leukemia is classified into M7. These methods of classification and of testing are known and are described in many textbooks (for example, Shin-Rinsho Naikagaku (New Clinical Internal Medicine), Fumimaro Takaku and Etsuro Ogata, Igakushoin Ltd., 1999).

35 [0007] Also, based on the identity of growing cells, chronic leukemia is roughly divided into chronic myelocytic leukemia (CML) and chronic lymphocytic leukemia (CLL). Also, as a subtype of chronic myelocytic leukemia, chronic myelomonocytic leukemia is known, and as a subtype of chronic lymphocytic leukemia, prolymphocytic leukemia is known.

40 [0008] Lymphoma is a generic term for tumors derived from cells constituting lymphatic tissues such as the lymph node, and is hematopoietic cell tumors caused mainly by canceration of lymphocytes. Malignant lymphoma is divided into Hodgkin's disease and non-Hodgkin lymphoma, both of which are cancerated lymphocytic cells and can be divided into the T lymphocytic and the B lymphocytic types.

45 [0009] As non-Hodgkin lymphoma, there are known B lymphocytic tumors such as follicular lymphoma, mantle cell lymphoma, Burkitt's lymphoma, pre-B lymphoma and the like. For T lymphocytic tumors, there are known adult T cell leukemia (ATL) and peripheral non-ATL T-lymphoma (PNTL). Also, diffuse lymphoma comprising two types (T lymphocytic and B lymphocytic) is included in non-Hodgkin lymphoma. Furthermore, as a subtype, hairy cell leukemia which is a B lymphocytic tumor is known.

50 [0010] Lymphocytic leukemia and lymphoma which are canceration of major constituent cells of lymphocytes are referred to as lymphocytic tumors, and are roughly divided into B lymphocytic tumors and T lymphocytic tumors. For example, B lymphocytic tumors include acute B lymphocytic leukemia (B-ALL), chronic B lymphocytic leukemia (B-CLL), pre-B lymphoma, Burkitt's lymphoma, follicular lymphoma, mantle cell lymphoma, diffuse lymphoma and the like. T lymphocytic tumors include acute T lymphocytic leukemia (T-ALL), chronic T lymphocytic leukemia (T-CLL), adult T cell leukemia (ATL), peripheral non-ATL T-lymphoma (PNTL) and the like (Zukai Rinshogan series (Illustrated Clinical Cancer Series) No. 17, Leukemia and lymphoma, Takashi Sugimura et al., MEDICAL VIEW Co., Ltd., 1987; B Lymphocytic Tumors, Kiyoshi Takatsuki, Nishimura Co., Ltd., 1991; Shin-Rinsho Naikagaku (New Clinical Internal

Medicine), Fumimaro Takaku and Etsuro Ogata, Igakushoin Ltd., 1999). Myeloma is also a type of lymphatic tumor, and exhibits characteristic clinical findings.

[0011] Myeloma which is also referred to as plasmacytoma and multiple myeloma is a neoplastic disease characterized by the accumulation of monoclonal plasma cells in the bone marrow. Myeloma is a disease in which plasma cells, i.e. terminally differentiated B cells that produce and secrete immunoglobulins, are monoclonally increased mainly in the bone marrow, and thus in the serum of patients with this disease, monoclonal immunoglobulins or components thereof, L chain, H chain, etc., can be detected.

[0012] For the treatment of myeloma, chemotherapeutic agents etc. have been used so far, but no effective therapeutic agents have been found that lead to the complete remission and the extension of the survival of patients. Thus, there has been a long-awaited need for agents having therapeutic effects based on a new mechanism of action. For lymphoma and leukemia as well, though moderately effective chemotherapy has been developed, new agents have been waited for due to adverse reactions.

[0013] On the other hand, Goto, T. et al. have reported a monoclonal antibody (mouse anti-HM1.24 antibody) that was obtained by immunizing mice with human plasma cells (Blood (1994) 84, 1922-1930). When anti-HM1.24 antibody was administered to a mouse transplanted with human myeloma cells, the antibody accumulated in tumor tissues in a specific manner (Masaaki Kosaka et al., Nippon Rinsho (Japan Clinical) (1995) 53, 627-635), suggesting that anti-HM1.24 antibody could be applied in the diagnosis of tumor localization by radioisotopic labeling, missile therapies such as radiotherapy, and the like.

[0014] In the above Blood (1994) 84, 1922-1930, it has been described that anti-HM1.24 antibody has an in vitro cytotoxicity on a human myeloma cell line RPMI8226. It has also been shown that chimeric anti-HM1.24 antibody, or anti-HM1.24 antibody that is mouse anti-HM1.24 antibody rendered chimeric, and a humanized reshaped anti-HM1.24 antibody, specifically bind to myeloma cells and have cytotoxicity (Blood (1999) 93, 3922-3920).

[0015] On the other hand, it has also been demonstrated for lymphocytic tumors that an antigen protein recognized by anti-HM1.24 antibody is expressed in lymphocytic tumors, and that anti-HM1.24 antibody has a cytotoxicity on lymphocytic tumors due to a CDC activity and an ADCC activity, and thereby exhibits anti-tumor effect (WO 98/35698). Thus, HM1.24 antigen has been highly expressed not only in myeloma cells that are terminally differentiated B cells but also in lymphocytic tumors, and anti-HM1.24 antibody that recognizes HM1.24 antigen is useful as a therapeutic agent for lymphocytic tumors.

[0016] Thus, HM1.24 antigen has been highly expressed in myeloma cells that are terminally differentiated B cells and in lymphocytic tumors, and anti-HM1.24 antibody that recognizes this antigen exhibits cell-killing activity in proportion to the number of HM1.24 antigens on the cell surface, and thus immunotherapy with anti-HM1.24 antibody is expected to provide an effective method of treating multiple myeloma and lymphocytic tumors. Thus, if the amount of HM1.24 antigen, which is an antigen against anti-HM1.2 antibody, expressed on the cell surface could be enhanced, the administration of a smaller amount of the antibody is expected to provide equivalent cytotoxicity, and it would become possible to decrease adverse reactions.

[0017] Furthermore, for hematopoietic tumor cells that are not expressing HM1.24 antigen, if the amount of HM1.24 antigen expressed on the cell surface could be enhanced, cytotoxicity or cytotoxic effect through ADCC activity or CDC activity with anti-HM1.24 antibody is expected for hematopoietic tumor cells for which, generally, anti-HM1.24 antibody alone is not effective.

[0018] On the other hand, interferon, that was discovered as a substance having an activity of inhibiting viral growth, is currently known to be classified into four groups of  $\alpha$ ,  $\beta$ ,  $\gamma$ , and  $\omega$  in mammals, and to have a variety of biological activities (Pestka, S., et al., Ann. Rev. Biochem. (1987) 56, 727-777; Langer, J. A., et al., Immunology Today (1988) 9, 393-400). However, there were no reports on whether interferon  $\alpha$  and interferon  $\gamma$  could have an effect of increasing the expressed amount of HM1.24 antigen in myeloma cells or cells of hematopoietic tumors such as lymphocytic tumors.

[0019] On the other hand, interferon regulatory factor (IRF)-1 and 2 were identified as a transcription regulatory factor of the IFN- $\beta$  gene. IRF-1 and 2 are generally known to bind to the same gene regulatory sequence, and act in an antagonistic manner in that IRF-1 acts as a transcription activation factor and IRF-2 as a transcription suppressing factor. The NIH3T3 cells in which IRF-2 was highly expressed has been demonstrated to exhibit enhanced cell saturation density, colony formation in the methylcellulose gel, and a tumorigenic property in nude mice, and IRF-2 has been shown to act as an oncogene.

[0020] On the other hand, recent advances in research have indicated that IRF-2 is required for the expression of histone H4 that acts in the control of cell cycle. IRF-2 has also been shown to increase the expression of vascular cell adhesion molecule-1 (VCAM-1) in muscle cells, and it has also been demonstrated that the acid region (182 to 218) of IRF-2 is involved in the activation of VCAM-1. Based on this, it is known that IRF-2 not only acts as a transcription regulatory factor but may act as a transcription activation factor.

[0021] However, it was not known that the IRF-2 protein binds to the promoter (HM1.24 promoter) of the HM1.24 antigen gene, and activates said promoter.

## DISCLOSURE OF THE INVENTION

[0022] Current methods of treating myeloma are, as mentioned above, not satisfactory and, accordingly, the appearance of epoch-making therapeutic drugs or methods that prolong the patient's survival is awaited. The treatment of myeloma with anti-HM1.24 antibody is likely to provide epoch-making therapeutic drugs in terms of specificity and effectiveness, and thus there is a need for methods of allowing anti-HM1.24 antibody to exhibit its effect more efficiently.

[0023] Also, as methods of treating lymphocytic tumors currently employed, there can be mentioned various chemotherapies, X-ray therapy, bone marrow transplantation and the like, but no therapeutic methods are satisfactory, and thus epoch-making therapeutic drugs or methods that attains complete remission of lymphocytic tumors and prolong the patient's survival are awaited. Furthermore, therapeutic methods that are effective for leukemia other than lymphocytic leukemia, i.e. myelocytic leukemia such as acute myelocytic leukemia and chronic lymphocytic leukemia are awaited. Therapeutic drugs and methods that attain complete remission of these myelomas, lymphocytic tumors, and myelocytic leukemia and that can prolong the patient's survival could provide therapeutic drugs and methods for hematopoietic tumors in general.

[0024] Thus, it is an object of the present invention to provide a means of enhancing the effect of anti-HM1.24 antibody of suppressing myeloma by enhancing the amount expressed of HM1.24 antigen in myeloma.

[0025] It is another object of the present invention to provide a means of enhancing the effect of anti-HM1.24 antibody of suppressing lymphocytic tumors by enhancing the amount expressed of HM1.24 antigen in lymphocytic tumors.

[0026] It is further object of the present invention to provide a means of inducing the effect of anti-HM1.24 antibody of suppressing myelocytic leukemia by providing a means of newly inducing the expression of HM1.24 antigen in myelocytic leukemia.

[0027] The above should provide a means of treating hematopoietic tumors with anti-HM1.24 antibody by enhancing the amount expressed of HM1.24 antigen or by using a means of newly inducing HM1.24 antigen.

[0028] After searching drugs that either enhance the amount expressed of HM1.24 antigen or allow HM1.24 antigen to be newly expressed on the cell surface, the present inventors have found that interferon  $\alpha$ , interferon  $\gamma$ , and the IRF-2 protein have the desired activity, and thereby have completed the present invention.

[0029] Thus the present invention provides an expression-enhancing or expression-inducing agent of a protein (HM1.24 antigen) having the amino acid sequence set forth in SEQ ID NO: 2 in hematopoietic tumor cells, said agent comprising interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein as an active ingredient.

[0030] The above hematopoietic tumors are, for example, leukemia, lymphoma, myeloma etc., and as this leukemia includes, for example, acute myelocytic leukemia, chronic myelocytic leukemia, acute lymphocytic leukemia, chronic lymphocytic leukemia etc., the above lymphoma includes, for example, Hodgkin's disease, T lymphocytic non-Hodgkin lymphoma, B lymphocytic non-Hodgkin lymphoma etc., and the above myeloma includes multiple myeloma.

[0031] The present inventors have also found that, by allowing an antigen that specifically binds to said HM1.24 antigen to bind to the hematopoietic tumor cells in which HM1.24 antigen has been expressed by the above expression-enhancing or expression-inducing agent of HM1.24 antigen, anti-tumor effect on said hematopoietic tumor cells can be enhanced.

[0032] Thus, the present invention also provides a therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors comprising, as an active ingredient, an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2, wherein interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein is used in combination.

[0033] The present invention also provides a therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors comprising, as an active ingredient, (1) interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein, and (2) an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.

[0034] The present invention also provides a therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors comprising, as an active ingredient, interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein, wherein an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2 is used in combination.

[0035] The above hematopoietic tumors are, for example, leukemia, lymphoma, or myeloma. The above leukemia includes, for example, acute myelocytic leukemia, chronic myelocytic leukemia, acute lymphocytic leukemia, chronic lymphocytic leukemia etc., the above lymphoma includes, for example, Hodgkin's disease, T lymphocytic non-Hodgkin lymphoma, B lymphocytic non-Hodgkin lymphoma etc., and the above myeloma includes multiple myeloma.

[0036] The above antibody is preferably an antibody having cytotoxicity, and the cytotoxicity is the ADCC activity. The antibody is preferably a monoclonal antibody, a chimeric antibody, a humanized antibody, or a human antibody.

The monoclonal antibody is preferably anti-HM1.24 antibody produced by a hybridoma having the Deposit No. FERM BP-5233, and said chimeric antibody or humanized antibody is preferably a chimeric antibody or a humanized antibody of anti-HM1.24 antibody produced by a hybridoma having the Deposit No. FERM BP-5233.

[0037] The present invention further provides a kit for the treatment of a patient having a hematopoietic tumor, said

kit comprising:

- (1) an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2; and
- (2) an instruction manual instructing the administration of the above antibody to the patient in combination with an expression-enhancing agent of a protein having the amino acid sequence set forth in SEQ ID NO: 2.

**[0038]** The present invention also provides a kit for the treatment of a patient having a hematopoietic tumor, said kit comprising:

- (1) an expression-enhancing agent of a protein having the amino acid sequence set forth in SEQ ID NO: 2; and
- (2) an instruction manual instructing the administration of the above antibody to the patient in combination with an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.

**[0039]** The present invention also provides a kit for the treatment of a patient having a hematopoietic tumor, said kit comprising:

- (1) an expression-enhancing agent of a protein having the amino acid sequence set forth in SEQ ID NO: 2; and
- (2) an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2; and
- (3) an instruction manual instructing the combined administration of the above agent and the above antibody to the patient.

**[0040]** The present invention also provides a method of screening an expression-enhancing agent of HM1.24 antigen, comprising the steps of:

- (1) preparing cells having a reporter gene that has the region of the HM1.24 gene promoter;
- (2) contacting said cells with a test substance; and
- (3) detecting the expression of the reporter gene.

**[0041]** The present invention also provides an expression-enhancing agent of HM1.24 antigen selected by the above method.

**[0042]** The present invention also provides a pharmaceutical composition for the treatment of hematopoietic tumors wherein said composition comprises the above expression-enhancing agent and said composition is used in combination with an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.

**[0043]** The present invention further provides a method of screening an expression-enhancing agent comprising the steps of:

- (1) contacting cells with a test substance;
- (2) determining the amount expressed of the IL-2 protein in said cells.

**[0044]** The present invention also provides an expression-enhancing agent of the IRF-2 protein selected by the above method.

**[0045]** The present invention also provides a pharmaceutical composition for the treatment of hematopoietic tumors wherein said composition comprises the above expression-enhancing agent and said composition is used in combination with an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.

#### BRIEF EXPLANATION OF THE DRAWINGS

**[0046]** Fig. 1 shows the result of an experiment in which a myeloma cell line U266 cultured in the absence (upper) or the presence (bottom) of interferon  $\alpha$  was analyzed by flow cytometry using human IgG (control) or anti-HM1.24 antibody as a label.

**[0047]** Fig. 2 shows the result of an experiment in which the myeloma cells of a patient cultured in the absence (upper) or the presence (bottom) of interferon  $\alpha$  was analyzed by flow cytometry using human IgG (control) or anti-HM1.24 antibody as a label.

**[0048]** Fig. 3 is a graph showing the result of an experiment in which U266 cells transformed with a reporter plasmid into which the promoter region of a gene encoding HM1.24 antigen had been inserted were cultured in the absence or in the presence of various concentrations of interferon  $\alpha$ , and then the luciferase activity was determined.

**[0049]** Fig. 4 is a graph showing the result of an experiment in which U266 cells or HEL cells transformed with a reporter plasmid into which a segment from the transcription initiation point to 151 bp upstream or to 77 bp upstream

among the promoter region of a gene encoding HM1.24 antigen had been inserted were cultured in the presence of interferon  $\alpha$  (1000 U/ml), and then the luciferase activity was determined.

[0050] Fig. 5 shows the result of an experiment in which a myeloma cell line U266 cultured in the absence (upper) or the presence (bottom) of interferon  $\gamma$  was analyzed by flow cytometry using human IgG (control) or anti-HM1.24 antibody as a label.

[0051] Fig. 6 shows the result of an experiment in which the myeloma cells of the patient cultured in the absence (upper) or the presence (bottom) of interferon  $\gamma$  were analyzed by flow cytometry using human IgG (control) or anti-HM1.24 antibody as a label.

[0052] Fig. 7 is an electrophoretogram that shows changes with time in the amount of a transcription factor that is produced by adding IFN  $\alpha$  to cultured U266 cells and that binds to the HM1.24 promoter region, and is a photograph substituting for  $\alpha$  drawing. NE(-): no nuclear extract was added. Oh: the nuclear extract without IFN- $\alpha$  stimulation was added. 0.5-8 h: the nuclear extracts that passed respective time after stimulation with IFN- $\alpha$  (1000 U/ml) was added. + cold: 50 ng of the nonlabeled ISRE2 probe was added. + cold unrelated: 50 ng of the nonlabeled adp sequence was added.

[0053] Fig. 8 is an electrophoretogram showing the result of an experiment in which a transcription factor that binds to HM1.24 promoter was identified using various antibodies, and is a photograph substituting for a drawing. NE(-): no nuclear extract was added. Oh: the nuclear extract without IFN- $\alpha$  stimulation was added. 8 h: the nuclear extract 8 hours after stimulation with IFN- $\alpha$  (1000 U/ml) was added. + cold: 50 ng of the nonlabeled ISRE2 probe was added. + cold unrelated: 50 ng of the nonlabeled adp sequence was added. Two  $\mu$ g of each antibody was added.

[0054] Fig. 9 is a graph showing the result of an experiment in which the HM1.24 promoter reporter plasmid and the IRF-2 expression plasmid were introduced into U266 cells, and the reporter activity was determined.

[0055] Fig. 10 shows the result of an experiment in which a human leukemia cell line HEL and cells collected from a patient with acute myelocytic leukemia were cultured in the absence (left) or the presence (right) of interferon  $\alpha$ , and then were analyzed by flow cytometry using human IgG (control) or anti-HM1.24 antibody as a label. The dark area represents the control and the light area represents the stain with anti-HM1.24 antibody.

[0056] Fig. 11 shows the result of an experiment in which cells collected from a patient with acute lymphocytic leukemia and a patient with B lymphocytic non-Hodgkin lymphoma were cultured in the absence (left) or the presence (right) of interferon  $\alpha$ , and then were analyzed by flow cytometry using human IgG (control) or anti-HM1.24 antibody as a label. The dark area represents the control and the light area represents the stain with anti-HM1.24 antibody.

[0057] Fig. 12 shows the result of an experiment in which cells collected from a patient with T lymphocytic non-Hodgkin lymphoma were cultured in the absence (left) or the presence (right) of interferon  $\alpha$ , and then were analyzed by flow cytometry using human IgG (control) or anti-HM1.24 antibody as a label. The dark area represents the control and the light area represents the stain with anti-HM1.24 antibody.

[0058] Fig. 13 shows the result of an experiment in which ADCC activity by various human anti-HM1.24 antibodies were determined when peripheral blood mononuclear cells from normal healthy subjects were used as the effector cell using a human leukemia cell line HEL as the standard cell.

## EMBODIMENTS FOR CARRYING OUT THE INVENTION

### Interferon- $\alpha$ and interferon- $\gamma$

[0059] Interferon was discovered as a substance having an activity of inhibiting viral growth and currently four types,  $\alpha$ ,  $\beta$ ,  $\gamma$ , and  $\omega$ , are known in mammals. In addition to the activity of inhibiting viral growth, they are known to exhibit an activity of inhibiting cell growth and of modulating immunological functions (Interferon "Cytokine", Toshiaki Osawa et al., (1990) 115-133, Tokyo Kagaku Dojin Co., Ltd.; Pestka, S., et al., Ann. Rev. Biochem. (1987) 56, 727-777; Langer, J. A., et al., Immunology Today (1988) 9, 393-400).

[0060] Interferon- $\alpha$  and interferon- $\gamma$  for use in the present invention may be mutants as long as they have an activity of increasing the amount expressed of HM1.24 antigen. In order to determine the amount expressed of HM1.24 antigen, as described in Examples, myeloma cells are harvested from a myeloma cell line or a patient with myeloma and then subjected to flow cytometry for detection. As mutants, they may be interferon- $\alpha$  and interferon- $\gamma$  in which one or several, or a plurality of amino acid residues have been deleted, substituted, or substituted or inserted.

[0061] As methods of introducing deletion, substitution, or insertion into proteins, there can be used site-directed mutagenesis that alters the corresponding gene (Hashimoto-Gotoh, Gene (1005) 152, 271-275, Zoller, Methods Enzymol. (1983) 100, 468-500, Kramer, Nucleic Acids Res. (1984) 12, 9441-8456, Kunkel, Proc. Natl. Acad. Sci. USA (1985) 82, 489-492, "New Cell Engineering Experimental Protocol" edited by Dept. of Oncology, Inst. of Medical Science, Univ. of Tokyo (1993) pp. 241-248).

[0062] It is also possible to use "Site-Directed Mutagenesis System" (GIBCO-BRL) and "QuickChange Site-Directed Mutagenesis Kit" (Stratagene) employing commercially available PCR. Amino acid mutations in proteins may some-

times take place in nature. That such a protein, in which mutation has been introduced, has an activity equal to the original protein has been shown in Mark, Proc. Natl. Acad. Sci. USA (1984) 81, 5662-5666.

**[0063]** In the substitution of amino acid residues, it is preferred to substitute between amino acids whose properties are conserved. For example, substitution is preferred between hydrophobic amino acids (A, I, L, M, F, P, W, Y, V), hydrophilic amino acids (R, D, N, C, E, Q, G, H, K, S, T), amino acids having aliphatic side chains (G, A, V, L, I, P), amino acids having hydroxyl group-containing side chains (S, T, Y), amino acids having sulfur-containing side chains (C, M), amino acids having carboxylic acid- and amide-containing side chains (D, N, E, Q), amino acids having base-containing side chains (R, K, H), and amino acids having aromatic group-containing side chains (H, F, Y, W).

**[0064]** Furthermore, as mutants, peptide fragments of interferon- $\alpha$  or interferon- $\gamma$  may be used. In particular, peptide fragments that have binding sites with interferon- $\alpha$  or interferon- $\gamma$  receptors are preferred. Preferably they are peptides comprised of 100 or more, more preferably 130 or more, still more preferably 150, and most preferably 160 or more contiguous amino acid residues.

#### IRF-2 protein

**[0065]** Interferon regulatory factors (IRF)-1 and -2 were identified as transcription regulatory factors of the IFN- $\beta$  gene (Taniguchi, T. et al., Nucleic Acids Res. (1989) 17, 8372; Taniguchi, T. et al., Cell (1989) 58, 729). IRF-1 and -2 are generally known to bind to the same gene regulatory sequence: IRF-1 and IRF-2 act in an antagonistic manner in that IRF-1 acts as a transcription activation factor, whereas IRF-2 as a transcription suppressing factor. The NIH3T3 cells in which IRF-2 is highly expressed has been demonstrated to exhibit enhanced cell saturation density, colony formation in the methylcellulose gel, and a tumorigenic property in nude mice, and IRF-2 acts as an oncogene.

**[0066]** On the other hand, recent advances in research have indicated that IRF-2 is required for the expression of histone H4 that acts for the control of cell cycle. IRF-2 is also shown to increase the expression of vascular cell adhesion molecule-1 (VCAM-1) in muscle cells, and it is becoming increasingly clear that the acid region (182 to 218) of IRF-2 is involved in the activation of VCAM-1. Based on this, it is known that IRF-2 not only acts as a transcription regulatory factor but as a transcription activation factor.

#### Hybridoma

**[0067]** The hybridoma that produces the antibody for use in the present invention can be basically constructed using a known technology as described below. Thus, the HM1.24 antigen protein or a HM1.24 antigen-expressing cell may be used as a sensitizing antigen and is immunized in the conventional method of immunization. The immune cells thus obtained are fused with known parent cells in the conventional cell fusion process, and then monoclonal antibody-producing cells are screened by the conventional screening method to construct the desired hybridoma.

**[0068]** Specifically, monoclonal antibody may be obtained in the following manner. For example, as the HM1.24 antigen-expressing cell which is the sensitizing antigen to obtain antibody, a human multiple myeloma cell line KPM2 (Japanese Unexamined Patent Publication (Kokai) No. 7-236475) and KPC-32 (Goto, T. et al., Jpn. J. Clin. Hematol. (1991) 32, 1400) can be used. As the sensitizing antigen, it is also possible to use a protein having the amino acid sequence set forth in SEQ ID NO: 1 or a peptide or a polypeptide containing an epitope recognized by anti-HM1.24 antibody.

**[0069]** The cDNA of the protein having the amino acid sequence set forth in SEQ ID NO 1 used as the sensitizing antigen may be inserted into the XbaI cleavage site of the pUC19 vector to prepare a plasmid pRS38-pUC19. E. coli having the plasmid pRS38-pUC19 has been internationally deposited under the provisions of the Budapest Treaty as Escherichia coli DH5 $\alpha$  (pRS38-pUC19) on October 5, 1993 with the Patent Microorganism Depository, the National Institute of Bioscience and Human Technology (Chuo Dai 6, 1-1, Higashi 1-chome, Tsukuba city, Ibaraki Pref., Japan) as FERM BP-4434 (Japanese Unexamined Patent Publication (Kokai) No. 7-196694). Using the cDNA fragment contained in this plasmid pRS38-pUC19, a peptide or a polypeptide that contains an epitope recognized by anti-HM1.24 antibody can be constructed by gene engineering technology.

**[0070]** Mammals to be immunized with the sensitizing antigen are not specifically limited, and they are preferably selected in consideration of their compatibility with the parent cell for use in cell fusion. They generally include rodents such as mice, rats, and hamsters.

**[0071]** Immunization of animals with a sensitizing antigen may be carried out using a known method. A general method, for example, involves the intraperitoneal or subcutaneous administration of a sensitizing antigen to the mammal.

**[0072]** Specifically, a sensitizing antigen which has been diluted and suspended in an appropriate amount of phosphate buffered saline (PBS) or physiological saline etc. is mixed, as desired, with an appropriate amount of a common adjuvant, for example Freund's complete adjuvant. After being emulsified, it is preferably administered to the mammal for several times every 4 to 21 days. Alternatively, a suitable carrier may be used at the time of immunization of the

sensitizing antigen.

[0073] After immunizing in this manner and confirming an increase in the desired antibody levels in the serum, immune cells are harvested from the mammal and are subjected to cell fusion. As immune cells to be subjected to cell fusion, there may be specifically mentioned spleen cells.

[0074] The mammalian myeloma cells as the other parent cells which are subjected to cell fusion with the above-mentioned immune cells preferably include various known cell lines such as P3X63Ag8.653 (J. Immunol. (1979) 123: 1548-1550), P3X63Ag8U.1 (Current Topics in Microbiology and Immunology (1978) 81: 1-7), NS-1 (Kohler, G. and Milstein, C., Eur. J. Immunol. (1976) 6: 511-519), MPC-11 (Margulies, D.H. et al., Cell (1976) 8: 405-415), SP2/0 (Shulman, M. et al., Nature (1978) 276: 269-270), FO (de St. Groth, S. F. et al. J. Immunol. Methods (1980) 35: 1-21), S194 (Trowbridge, I. S., J. Exp. Med. (1978) 148: 313-323), R210 (Galfre, G. et al., Nature (1979) 277: 131-133) and the like.

[0075] Cell fusion between the above immune cells and the myeloma cells may be essentially conducted in accordance with a known method such as is described in Milstein et al. (Kohler, G. and Milstein, C., Methods Enzymol. (1981) 73: 3-46) and the like.

[0076] More specifically, the above cell fusion is carried out in the conventional nutrient broth in the presence of, for example, a cell fusion accelerator. As the cell fusion accelerator, for example, polyethylene glycol (PEG), Sendai virus (HVJ) and the like may be used, and, in addition, an adjuvant such as dimethyl sulfoxide etc. may be added as desired to enhance the efficiency of the fusion.

[0077] The preferred ratio of the immune cells and the myeloma cells to be used is, for example, 1 to 10 times more immune cells than the myeloma cells. Examples of culture media to be used for the above cell fusion include RPMI1640 medium and MEM culture medium suitable for the growth of the above myeloma cell lines, and the conventional culture medium used for this type of cell culture, and besides a serum supplement such as fetal calf serum (FCS) may be added.

[0078] In cell fusion, predetermined amounts of the above immune cells and the myeloma cells are mixed well in the above culture liquid, to which a PEG solution previously heated to about 37 °C, for example a PEG solution with a mean molecular weight of about 1000 to 6000, is added at a concentration of 30 to 60% (w/v), and mixed to obtain the desired fusion cells (hybridomas). Then by repeating the sequential addition of a suitable culture liquid and centrifugation to remove the supernatant, cell fusion agents etc. which are undesirable for the growth of the hybridoma can be removed.

[0079] Said hybridoma is selected by culturing in a conventional selection medium, for example, the HAT culture medium (a culture liquid containing hypoxanthine, aminopterin, and thymidine). Culturing in said HAT culture medium is continued generally for a period of time sufficient to effect killing of the cells (non-fusion cells) other than the desired hybridoma, generally several days to several weeks. Then, the conventional limiting dilution method is conducted in which the hybridomas that produce the desired antibody are screened and monoclonally cloned.

[0080] In addition to obtaining the above hybridoma by immunizing an animal other than the human with an antigen, it is also possible to sensitize human lymphocytes in vitro with HM1.24 antigen or HM1.24 antigen-expressing cells, and to allow the resulting sensitized lymphocytes to be fused with a human myeloma cell, for example U266, and thereby to obtain the desired human antibody having the activity of binding to HM1.24 antigen or to HM1.24 antigen-expressing cells (see Japanese Examined Patent Publication (Kokoku) No. 1-59878). Furthermore, a transgenic animal having a repertoire of human antibody genes is immunized with HM1.24 antigen or HM1.24 antigen-expressing cells to obtain the desired antibody according to the above-mentioned method (see International Patent Publication WO 93/12227, WO 92/03918, WO 94/02602, WO 94/25585, WO 96/34096, and WO 96/33735).

[0081] Furthermore, using a human antibody library, the desired human antibody may be isolated by means of panning. For example, the variable region of human antibody is expressed on the surface of a phage by the phage display method as a single chain antibody (scFv) to select a phage that binds to the HM1.24 antigen using a HM1.24 antigen-immobilized plate. By analyzing the gene of the phage selected, the gene encoding the variable region of the human antibody that binds to the HM1.24 antigen can be identified. Using these gene sequences, anti-HM1.24 antibody can be prepared. These methods are already known and can be found in WO 92/01047, WO 92/20791, WO 93/06213, WO 93/11236, WO 93/19172, WO 95/01438, and WO 95/15388.

[0082] The monoclonal antibody-producing hybridomas thus constructed can be subcultured in the conventional culture liquid, or can be stored for a prolonged period of time in liquid nitrogen.

[0083] In order to obtain monoclonal antibodies from said hybridoma, there may be employed a method in which said hybridoma is cultured by the conventional method and the antibodies are obtained as the culture supernatant, or a method in which the hybridoma is administered to and grown in a mammal compatible with said hybridoma and the antibodies are obtained as the ascites, or other methods. The former method is suitable for obtaining high-purity antibodies, whereas the latter is suitable for a large scale production of antibodies.

#### Monoclonal antibody

[0084] Specifically the anti-HM1.24 antibody-producing hybridoma can be constructed using the method of Goto, T.



et al. (Blood (1994) 84: 1922-1930). It can be conducted by: a method in which the anti-HM1.24 antibody-producing hybridoma that was internationally deposited under the provisions of the Budapest Treaty as FERM BP-5233 on April 27, 1995 with the Patent Microorganism Depository, the National Institute of Bioscience and Human Technology (Chuo Dai 6, 1-1, Higashi 1-chome, Tsukuba city, Ibaraki Pref., Japan) is intraperitoneally injected to BALB/c mice (manufactured by CLEA Japan) to obtain the ascites, from which the anti-HM1.24 antibody is purified, or: a method in which said hybridoma is cultured in a suitable culture medium such as the RPMI1640 medium containing 10% bovine fetal serum and 5% BM-Condensed H1 (manufactured by Boehringer Mannheim), the hybridoma SFM medium (manufactured by GIBCO-BRL), the PFHM-II medium (manufactured by GIBCO-BRL) and the like, and the anti-HM1.24 antibody can be purified from the supernatant.

#### Recombinant antibody

**[0085]** A recombinant antibody which was produced by the recombinant gene technology, in which an antibody gene was cloned from the hybridoma and integrated into a suitable vector which was then introduced into a host, can be used in the present invention as monoclonal antibody (see, for example, Carl, A. K., Borrebaeck, and James, W. Larrick, THERAPEUTIC MONOCLONAL ANTIBODIES, published in the United Kingdom by MACMILLAN PUBLISHERS LTD. 1990).

**[0086]** Specifically, mRNA encoding the variable region (V) of the desired antibody is isolated from the hybridoma producing the antibody. The isolation of mRNA is conducted by preparing total RNA using, for example, a known method such as the guanidine ultracentrifuge method (Chirgwin, J.M. et al., Biochemistry (1979) 18, 5294-5299), the AGPC method (Chmczynski, P. et al., (1987) 162, 156-159), and then mRNA is purified from the total RNA using the mRNA Purification kit (manufactured by Pharmacia) and the like. Alternatively, mRNA can be directly prepared using the QuickPrep mRNA Purification Kit (manufactured by Pharmacia).

**[0087]** cDNA of the V region of the antibody may be synthesized using a reverse transcriptase from the mRNA thus obtained. cDNA may be synthesized using the AMV Reverse Transcriptase First-strand cDNA Synthesis Kit and the like. Alternatively, for the synthesis and amplification of cDNA, the 5'-Ampli FINDER RACE Kit (manufactured by Clontech) and the 5'-RACE method (Frohman, M.A. et al., Proc. Natl. Acad. Sci. U.S.A. (1988) 85, 8998-9002; Belyavsky, A. et al., Nucleic Acids Res. (1989) 17, 2919-2932) that employs PCR may be used. The desired DNA fragment is purified from the PCR product obtained and may be ligated to vector DNA. Moreover, a recombinant vector is constructed therefrom and then is introduced into E. coli etc., from which colonies are selected to prepare the desired recombinant vector. The base sequence of the desired DNA may be confirmed by a known method such as the dideoxy method.

**[0088]** Once the DNA encoding the V region of the desired antibody has been obtained, it may be ligated to DNA encoding the constant region (C region) of the desired antibody, which is then integrated into an expression vector. Alternatively, the DNA encoding the V region of the antibody may be integrated into an expression vector which already contains DNA encoding the C region of the antibody.

**[0089]** In order to produce the antibody for use in the present invention, the antibody gene is integrated as described below into an expression vector so as to be expressed under the control of the expression regulatory region, for example an enhancer and/or a promoter. Subsequently, the expression vector may be transformed into a host cell and the antibody can then be expressed therein.

#### Altered antibody

**[0090]** In accordance with the present invention, artificially altered recombinant antibody such as chimeric antibody and humanized antibody can be used for the purpose of lowering heterologous antigenicity against humans. These altered antibody can be produced using known methods.

**[0091]** Chimeric antibody can be obtained by ligating the thus obtained DNA encoding the V region of antibody to DNA encoding the C region of human antibody, which is then integrated into an expression vector and introduced into a host for production of the antibody therein (see European Patent Application EP 125023, and International Patent Publication WO 96/02576). Using this known method, chimeric antibody useful for the present invention can be obtained.

**[0092]** For example, E. coli having the plasmid that contains the L chain and the H chain of chimeric anti-HM1.24 antibody has been internationally deposited under the provisions of the Budapest Treaty as Escherichia coli DH5 $\alpha$  (pUC19-1.24L-gK) and Escherichia coli DH5 $\alpha$  (pUC19-1.24H-gY1), respectively, on August 29, 1996 with the Patent Microorganism Depository, the National Institute of Bioscience and Human Technology (Chuo Dai 6, 1-1, Higashi 1-chome, Tsukuba city, Ibaraki Pref., Japan) as FERM BP-5646 and FERM BP-5644, respectively (see Japanese Patent Application No. 8-264756).

**[0093]** Humanized antibody which is also referred to as reshaped human antibody has been made by transplanting

the complementarity determining region (CDR) of antibody of a mammal other than the human, for example mouse antibody, into the CDR of human antibody. The general recombinant DNA technology for preparation of such antibodies is also known (see European Patent Application EP 125023 and International Patent Publication WO 96/02576).

[0094] Specifically, a DNA sequence which was designed to ligate the CDR of mouse antibody with the framework region (FR) of human antibody is synthesized by the PCR method from several divided oligonucleotides having sections overlapping with one another at the ends thereof. The DNA thus obtained is ligated to the DNA encoding the C region of human antibody, and then is integrated into an expression vector, which is introduced into a host for antibody production (see European Patent Application EP 239400 and International Patent Publication WO 96/02576).

[0095] For the FR of human antibody ligated through CDR, the FR is selected for which the complementarity determining region forms a favorable antigen binding site. When desired, amino acids in the framework region of the antibody variable region may be substituted so that the complementarity determining region of reshaped human antibody may form an appropriate antigen binding site (Sato, K. et al., Cancer Res. (1993) 53, 851-856).

[0096] For example, E. coli having the plasmid that contains the L chain and the H chain of humanized anti-HM1.24 antibody has been internationally deposited under the provisions of the Budapest Treaty as Escherichia coli DH5 $\alpha$  (pUC19-RVLa-AHM-gk) and Escherichia coli DH5 $\alpha$  (pUC19-RVHr-AHM-gy1), respectively, on August 29, 1996 with the Patent Microorganism Depository, the National Institute of Bioscience and Human Technology (Chuo Dai 6, 1-1, Higashi 1-chome, Tsukuba city, Ibaraki Pref., Japan) as FERM BP-5645 and FERM BP-5643, respectively (Patent Application No. 8-264756).

[0097] For chimeric antibody or humanized antibody, the C region of human antibody is used and, as the C region of human antibody that exhibits cytotoxicity, human C $\gamma$ , for example C $\gamma$ 1, C $\gamma$ 2, C $\gamma$ 3, and C $\gamma$ 4, can be used. Among them, antibody having C $\gamma$ 1 and C $\gamma$ 3 in particular has potent cytotoxicity, i.e. ADCC activity and CDC activity, and is used preferably in the present invention.

[0098] Chimeric antibody consists of the variable region of antibody derived from a mammal other than the human and the C region derived from human antibody, whereas humanized antibody consists of the complementarity determining region of antibody derived from a mammal other than the human and the framework region (FR) and the C region of antibody derived from human antibody. Accordingly, antigenicity thereof in the human body has been reduced so that they are useful as the active ingredient of the therapeutic agents of the present invention.

[0099] A preferred embodiment of the humanized antibody for use in the present invention includes humanized anti-HM1.24 antibody (see WO 98/14580).

### Expression and production

[0100] Antibody genes constructed as described above may be expressed and obtained in a known method. In the case of mammalian cells, expression may be accomplished using an expression vector containing a commonly used useful promoter, the antibody gene to be expressed, and DNA in which the poly A signal has been operably linked at 3' downstream thereof or a vector containing said DNA. Examples of the promoter/enhancer include human cytomegalovirus immediate early promoter/enhancer.

[0101] Additionally, as the promoter/enhancer which can be used for the expression of antibody for use in the present invention, there can be used viral promoters/enhancers such as retrovirus, polyoma virus, adenovirus, and simian virus 40 (SV40), and promoters/enhancers derived from mammalian cells such as human elongation factor 1 $\alpha$  (HEF1 $\alpha$ ).

[0102] For example, expression may be readily accomplished by the method of Mulligan et al. (Nature (1979) 277, 108) when SV40 promoter/enhancer is used, or by the method of Mizushima et al. (Nucleic Acids Res. (1990) 18, 5322) when HEF1 $\alpha$  promoter/enhancer is used.

[0103] In the case of E. coli, expression may be conducted by operably linking a commonly used useful promoter, a signal sequence for antibody secretion, and the antibody gene to be expressed, followed by expression thereof. As the promoter, for example, there can be mentioned lacZ promoter and araB promoter. The method of Ward et al. (Nature (1998) 341, 544-546; FASEB J. (1992) 6, 2422-2427) may be used when lacZ promoter is used, and the method of Better et al. (Science (1988) 240, 1041-1043) may be used when araB promoter is used.

[0104] As the signal sequence for antibody secretion, when produced in the periplasm of E. coli, the pelB signal sequence (Lei, S.P. et al., J. Bacteriol. (1987) 169, 4379) can be used. After separating the antibody produced in the periplasm, the structure of the antibody is appropriately refolded before use (see, for example, WO 96/30394).

[0105] As the origin of replication, there can be used those derived from SV40, polyoma virus, adenovirus, bovine papilloma virus (BPV) and the like. Furthermore, for the amplification of the gene copy number in the host cell system, expression vectors can include as selectable markers the aminoglycoside transferase (APH) gene, the thymidine kinase (TK) gene, E. coli xanthine guaninephosphoribosyl transferase (Ecogpt) gene, the dihydrofolate reductase (dhfr) gene and the like.

[0106] For the production of antibody for use in the present invention, any production system can be used. The production system of antibody preparation comprises the in vitro and the in vivo production system. As the in vitro

production system, there can be mentioned a production system which employs eukaryotic cells and the production system which employs prokaryotic cells.

[0107] When the eukaryotic cells are used, there are the production systems which employ animal cells, plant cells, and fungal cells. Known animal cells include (1) mammalian cells such as CHO cells, COS cells, myeloma cells, baby hamster kidney (BHK) cells, HeLa cells, and Vero cells, (2) amphibian cells such as *Xenopus* oocytes, or (3) insect cells such as sf9, sf21, and Tn5. Known plant cells include, for example, those derived from the genus *Nicotiana*, more specifically cells derived from *Nicotiana tabacum*, which is subjected to callus culture. Known fungal cells include yeasts such as the genus *Saccharomyces*, more specifically *Saccharomyces cereviceae*, or filamentous fungi such as the genus *Aspergillus*, more specifically *Aspergillus niger*.

[0108] When the prokaryotic cells are used, there are the production systems which employ bacterial cells. Known bacterial cells include *Escherichia coli* (*E. coli*), and *Bacillus subtilis*.

[0109] By introducing, via transformation, the gene of the desired antibody into these cells and culturing the transformed cells in vitro, the antibody can be obtained. Culturing is conducted in the known methods. For example, as the culture liquid, DMEM, MEM, RPMI1640, and IMDM can be used, and serum supplements such as fetal calf serum (FCS) may be used in combination. In addition, antibodies may be produced in vivo by implanting cells into which the antibody gene has been introduced into the abdominal cavity of an animal and the like.

[0110] As further in vivo production systems, there can be mentioned those which employ animals and those which employ plants. When animals are used, there are the production systems which employ mammals and insects.

[0111] As mammals, goats, pigs, sheep, mice, and cattle can be used (Vicki Glaser, SPECTRUM Biotechnology Applications, 1993). As insects, silkworms can be used.

[0112] When plants are used, tobacco, for example, can be used.

[0113] Antibody genes are introduced into these animals or plants, and the antibodies are produced in such animals or plants, and recovered therefrom. For example, an antibody gene is inserted into the middle of the gene encoding protein which is inherently produced in the milk such as goat  $\beta$  casein to prepare fusion genes. DNA fragments containing the fusion gene into which the antibody gene has been inserted are injected into a goat embryo, and the embryo is introduced into a female goat. The desired antibody is obtained from the milk produced by the transgenic goat born to the goat who received the embryo or offsprings thereof. In order to increase the amount of milk containing the desired antibody produced by the transgenic goat, hormones may be given to the transgenic goat as appropriate. (Ebert, K. M. et al., *Bio/Technology* (1994) 12, 699-702).

[0114] When silkworms are used, baculovirus into which the desired antibody gene has been inserted is infected to the silkworm, and the desired antibody can be obtained from the body fluid of the silkworm (Susumu, M. et al., *Nature* (1985) 315, 592-594). Moreover, when tobacco is used, the desired antibody gene is inserted into an expression vector for plants, for example pMON 530, and then the vector is introduced into a bacterium such as *Agrobacterium tumefaciens*. The bacterium is then infected to tobacco such as *Nicotiana tabacum* to obtain the desired antibody from the leaves of the tobacco (Julian, K.-C. Ma et al., *Eur. J. Immunol.* (1994) 24, 131-138).

[0115] When antibody is produced in the in vitro or in vivo production systems, as described above, DNA encoding the heavy chain (H chain) or the light chain (L chain) of antibody may be separately integrated into an expression vector and the hosts are transformed simultaneously, or DNA encoding the H chain and the L chain may be integrated into a single expression vector and the host is transformed therewith (see International Patent Publication WO 94-11523).

[0116] The antibody produced as described above can be bound to various molecules such as polyethylene glycol (PEG) for use as a modified antibody. "Antibody" as used herein includes these modified antibodies. In order to obtain these modified antibody, the antibody obtained may be chemically modified. These methods have already been established in the field of the art.

#### Separation and purification of antibody

[0117] Antibodies produced and expressed as described above can be separated from the inside or outside of the cell or from the host and then may be purified to homogeneity. Separation and purification of the antibody for use in the present invention may be accomplished by affinity chromatography. As the column used for such affinity chromatography, there can be mentioned Protein A column and Protein G column. Examples of the column employing Protein A column are Hyper D, POROS, Sepharose F.F. and the like.

[0118] Alternatively, methods for separation and purification conventionally used for proteins can be used without any limitation. Separation and purification of the antibody for use in the present invention may be accomplished by combining, as appropriate, chromatography other than the above-mentioned affinity chromatography, filtration, ultra-filtration, salting-out, dialysis and the like. Chromatography includes, for example, ion exchange chromatography, hydrophobic chromatography, gel-filtration and the like.

Determination of antibody concentration

[0119] The concentration of antibody obtained in the above method can be determined by the measurement of absorbance or by ELISA and the like. Thus, when absorbance measurement is employed, the antibody for use in the present invention or a sample containing the antibody is appropriately diluted with PBS(-) and then the absorbance is measured at 280 nm, followed by calculation using the absorption coefficient of 1.35 OD at 1 mg/ml. When the ELISA method is used, measurement is conducted as follows. Thus, 100  $\mu$ l of goat anti-human IgG (manufactured by BIO SOURCE) diluted to 1  $\mu$ g/ml in 0.1 M bicarbonate buffer, pH 9.6, is added to a 96-well plate (manufactured by Nunc), and is incubated overnight at 4 °C to immobilize the antibody.

[0120] After blocking, 100  $\mu$ l each of appropriately diluted antibody of the present invention or a sample containing the antibody, or 100  $\mu$ l of human IgG (manufactured by CAPPEL) as the standard is added, and incubated at room temperature for 1 hour. After washing, 100  $\mu$ l of 5000-fold diluted alkaline phosphatase-labeled anti-human IgG antibody (manufactured by BIO SOURCE) is added, and incubated at room temperature for 1 hour. After washing, the substrate solution is added and incubated, followed by the measurement of absorbance at 405 nm using the MICROPLATE READER Model 3550 (manufactured by Bio-Rad) to calculate the concentration of the desired antibody.

FCM analysis

[0121] Reactivity of the antibody of the present invention with lymphocytes may be examined by flow cytometry (FCM) analysis. As the cells, established cell lines or freshly isolated cells can be used. As established cell lines, there may be used myeloma-derived RPMI8226 (ATCC CCL 155), myeloma-derived U266 (ATCC TIB 196), myeloma-derived KPMM2, myeloma-derived KPC-32, and plasmacytoma-derived ARH-77 (ATCC CRL 1621), and the like.

[0122] After washing the above cells in PBS(-), 100  $\mu$ l of antibody or a control antibody diluted to 25  $\mu$ g/ml in the FACS buffer (PBS(-) containing 2% bovine fetal serum and 0.05% sodium azide) is added thereto, which is then incubated on ice for 30 minutes. After washing with the FACS buffer, 100  $\mu$ l of 25  $\mu$ g/ml FITC-labeled goat anti-mouse antibody (GAM, manufactured by Becton Dickinson) is added thereto, which is then incubated on ice for 30 minutes. After washing with the FACS buffer, the cells are suspended in 600  $\mu$ l of 1 ml of the FACS buffer, and each cell may be measured for its fluorescence intensity using the FACScan (manufactured by Becton Dickinson).

Screening method

[0123] In order to screen the expression enhancer of HM1.24 antigen, for example, the cells that have not been stimulated and that are not expressing HM1.24 antigen or the cells that at least are expressing the antigen are determined using FCM analysis. For example, the cells described in Examples and a test substance are incubated for 1-2 days, and then stained with mouse anti-human HM1.24 antibody as a primary antibody. The cells are washed and further stained with FITC-labeled anti-mouse IgG antibody as a secondary antibody. Finally, after washing the cells, the fluorescence intensity of FITC is measured by a flow cytometer.

[0124] Furthermore, instead of the above indirect staining, FCM analysis by direct staining may be used in which the cells are treated with a high concentration of immunoglobulin, and then stained, after blocking Fc receptors, with FITC-labeled anti-human HM1.24 antibody.

[0125] It is also possible to screen expression enhancers of HM1.24 antigen by the reporter gene assay using the HM1.24 promoter sequence. As the reporter gene, luciferase can be used. A plasmid is constructed that contains the HM1.24 promoter sequence upstream of the reporter gene, after which it is transformed into the cells, and the cells obtained are cultured with a test substance for 1-2 days, and the cells recovered are subjected to FCM analysis to screen drugs that enhance the expression of HM1.24 antigen.

CytotoxicityMeasurement of ADCC activity

[0126] The antibody for use in the present invention is one which has, for example, an ADCC activity as the cytotoxicity.

[0127] According to the present invention, the ADCC activity on hematopoietic tumor cells can be measured in the following manner. First, mononuclear cells (E) are isolated as the effector cells from human peripheral blood or bone marrow by the gravity centrifuge method.

[0128] As the target cells (T), RPMI8226 (ATCC CCL 155), U266 (ATCC TIB 196), KPMM2, KPC-32, ARH-77 (ATCC CRL 1621), HEL, cells derived from patients or the like is labeled with <sup>51</sup>Cr to be prepared as the target cells. Subsequently, to the labeled target cells is added the antibody to be measured for the ADCC activity and incubated. Effector

cells at a suitable ratio to the target cells are then added and incubated.

**[0129]** After incubation, the supernatant is removed and measured for radioactivity using a gamma counter. At this time 1% NP-40 can be used for measurement of the maximum free radioactivity. The cytotoxicity (%) can be calculated as  $(A-C)/(B-C) \times 100$ , in which A is radioactivity (cpm) liberated in the presence of the antibody, B is radioactivity (cpm) liberated by NP-40, and C is radioactivity (cpm) liberated by the medium alone containing no antibody.

#### Enhancement of cytotoxicity

**[0130]** In order to exhibit cytotoxicity such as an ADCC activity, it is preferred to use  $C\gamma$ , in particular  $C\gamma 1$  and  $C\gamma 3$  as the constant region (C region) of antibody in humans. Furthermore, a more potent ADCC activity or CDC activity can be induced by adding, altering, or modifying part of the amino acids in the C region of antibody.

**[0131]** By way of example, there can be mentioned the conversion of IgG to an IgM-like polymer by amino acid substitution (Smith, R.I.F. & Morrison, S.L. BIO/TECHNOLOGY (1994) 12, 683-688), the conversion of IgG to an IgM-like polymer by amino acid addition (Smith, R.I.F. et al., J. Immunology (1995) 154, 2226-2236), the expression of a tandemly-ligated gene encoding L chain (Shuford, W. et al., Science (1991) 252, 724-727), the dimerization of IgG by amino acid substitution (Caron, P.C. et al., J. Exp. Med. (1992) 176, 1191-1195, Shopes, B., J. Immunology (1992) 148, 2918-2922), the dimerization of IgG by chemical modification (Wolff, E. A. et al., Cancer Res. (1993) 53, 2560-2565), and the introduction of the effector function by altering an amino acid(s) in the hinge region of antibody (Norderhaug, L. et al., Eur. J. Immunol. (1991) 21, 2379-2384) and the like.

**[0132]** These can be accomplished by means of the oligomer site-specific mutagenesis using a primer, the addition of a base sequence using a restriction enzyme cleavage site, and the use of a chemical modifier that creates a covalent bond.

#### Treatment of patients

**[0133]** One embodiment of the present invention concerns a method of treating hematopoietic tumors by administering to the patient a pharmaceutical agent that enhances the amount expressed of HM1.24 antigen, and a pharmaceutical agent that induces the expression of HM1.24 antigen in the cell that normally is not expressing HM1.24 antigen on the cell surface and anti-HM1.24 antibody. The hematopoietic tumors are, for example, myeloma, preferably multiple myeloma, lymphocytic tumors such as lymphoma, preferably Hodgkin's disease or non-Hodgkin lymphoma, lymphocytic leukemia, preferably acute T lymphocytic leukemia, chronic T lymphocytic leukemia, acute B lymphocytic leukemia, chronic B lymphocytic leukemia, and myelocytic leukemia for example acute myelocytic leukemia and chronic myelocytic leukemia. Furthermore, acute leukemia belonging to L1-L3 and M0-M7 in the FAB Classification is also included.

**[0134]** A pharmaceutical agent that enhances the amount expressed of HM1.24 antigen, and a pharmaceutical agent that induces the expression of HM1.24 antigen in the cell that normally is not expressing HM1.24 antigen on the cell surface and anti-HM1.24 antibody is preferably interferon- $\alpha$ , interferon- $\gamma$ , the IRF-2 protein, or a vector containing DNA encoding the IRF-2 protein, and preferably interferon- $\alpha$  and interferon- $\gamma$ . The dosage of interferon- $\alpha$  and interferon- $\gamma$  is, when administered to humans, is an amount that attains the highest blood level of 1-10000 I.U./ml (International Units), more preferably 5-1000 I.U./ml, even more preferably 5-500 I.U./ml, and most preferably 5-50 I.U./ml.

**[0135]** In the case of intravenous administration, preferably ten thousand to ten million I.U., more preferably one hundred thousand to ten million I.U., even more preferably fifty thousand to five million I.U., most preferably one million to five million I.U. is given per administration. Interferon and anti-HM1.24 antibody may be administered together or separately. In the latter case, preferably interferon is given first, followed by the administration of anti-HM1.24 antibody within 96 hours. The interval between the interferon administration and the anti-HM1.24 antibody administration is not limited as long as the amount expressed of HM1.24 antigen is being enhanced by the administration of interferon, but it is preferably within 96 hours, more preferably 72 hours, still more preferably 48 hours.

**[0136]** Alternate administration of interferon and anti-HM1.24 antibody for a plurality of times depending on the clinical response of the patient is within the scope of the present invention. The route of administration is preferably given directly into the blood circulation, and intravenous administration or intraarterial administration is preferred. Continued administration is possible and intravenous drip may be used. The administration may also be subcutaneous or intramuscular administration.

**[0137]** Another aspect of the present invention concerns a therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors comprising interferon- $\alpha$  or interferon- $\gamma$  and anti-HM1.24 antibody. The therapeutic agent of the present invention may contain a pharmaceutically acceptable vehicle that has been used for interferon and antibody preparations, such as physiological saline or 5% dextran, together with a common stabilizer or a excipient.

**[0138]** Another aspect of the present invention provides a kit for treating a patient with hematopoietic tumors, com-

prising a pharmaceutical composition comprising anti-HM1.24 antibody as an active ingredient and an instruction manual that contains description of combined therapy with interferon- $\alpha$  or interferon- $\gamma$ .

**[0139]** Another aspect of the present invention provides a pharmaceutical composition comprising anti-HM1.24 antibody as an active ingredient for treating a patient with hematopoietic tumors, wherein said composition is used in combination with interferon- $\alpha$  or interferon- $\gamma$ .

**[0140]** Another aspect of the present invention relates to the administration of a gene encoding a substance that enhances or induces the expression of HM1.24 antigen. For example, the DNA sequences of interferon- $\alpha$ , interferon- $\gamma$ , IRF-2 are known and can be administered to patients by integration into the desired vector. As the vector, they are inserted into adenovirus vector (for example pAdexLew) or retrovirus vector (for example pZIPneo), which are then administered to the living body. The method of administration may be ex vivo or in vivo. Alternatively, naked DNA may be administered.

## EXAMPLES

### Example 1. Enhancement of the amount of HM1.24 antigen expressed in myeloma cells by interferon- $\alpha$

**[0141]** A human myeloma cell line U266 (ATCC TIB 196) and myeloma cells derived from the bone marrow of a patient with multiple myeloma were cultured in a RPMI1640 medium (Sigma, St Louis, MO, USA) containing 10% fetal bovine serum (Whittaker Bioproducts, Inc., Walkersville, MD, USA) in a 5% carbon dioxide incubator at 37°C. The hybridoma that produces mouse anti-HM1.24 antibody has been internationally deposited as FERM BP-5233 (deposition date: April 27, 1995) with the National Institute of Bioscience and Human Technology, of 1-3, Higashi 1-chome, Tsukuba city, Ibaraki Pref.

**[0142]** Myeloma cells ( $1 \times 10^5$ /ml) were cultured in the presence or absence of 1000 U/ml of the natural type interferon- $\alpha$  (Otsuka Pharmaceutical, Tokyo) for 48 hours, and changes in HM1.24 antigen (the base sequence encoding this is shown in SEQ ID NO: 1) were determined by flow cytometry. After the cells were washed with phosphate buffer (Gibco BRL, Grand Island, NY, USA) supplemented with 0.1% bovine serum albumin (Sigma, St Louis, MO, USA) and 0.02% sodium azide, they were suspended into PBS (100  $\mu$ l) supplemented with human immunoglobulin (3 mg/ml, Green Cross, Osaka), and were allowed to react at 4 °C for 15 minutes.

**[0143]** Thereafter, 2  $\mu$ l of FITC-human IgG1 (1 mg/ml) or FITC-anti-HM1.24 antibody (1 mg/ml) was added to stain at 4 °C for 60 minutes. When the patient's myeloma cells were used, 20  $\mu$ l of PE-anti-CD38 (Becton Dickinson, San Jose, CA, USA) was added for double staining to identify the myeloma cells. After staining, the cells were washed twice with PBC, and were stored in PBS containing 1% paraformaldehyde (Wako Pure Chemical Industries, Ltd., Osaka). Subsequently, the expression of HM1.24 antigen was analyzed using a flow cytometer (EPICS XL, Coulter, Hi-aleah, FL, USA).

**[0144]** As a result, a myeloma cell line U266 (Figure 1) and the patient's myeloma cells (Figure 2) were expressing HM1.24 antigen at a condition of no stimulation, and the stimulation with interferon- $\alpha$  further increased the amount expressed of HM1.24 antigen.

**[0145]** Interferon- $\alpha$  further enhanced the expression of HM1.24 antigen in the myeloma cell, and increased the number of anti-HM1.24 antibodies that bind to the myeloma cell. Since the therapeutic anti-tumor effect by anti-HM1.24 antibody is proportional to the number of antibodies that bound, treatment with anti-HM1.24 antibody after the administration of interferon- $\alpha$  is expected to provide a therapy that enhances the therapeutic effect by antibody and further enhances effectiveness.

### Example 2. Analysis of the expression function of HM1.24 antigen by the reporter gene analysis

**[0146]** In order to investigate whether the expression induction of antigen is regulated by the HM1.24 promoter region, the reporter gene at the promoter region was analyzed.

**[0147]** The gene (SEQ ID NO: 3) of the HM1.24 promoter region was obtained by PCR cloning. Genomic DNA was prepared from human peripheral blood mononuclear cells using the DNAzol reagent (GIBCO). With the genomic DNA obtained as the template, using primer HM2k (aaagggtaccagctgtcttctgtctgtcc) (SEQ ID NO: 4) and BST2B (atagtcatacgaagtagatgccatccag) (SEQ ID NO: 5), PCR (94°C for 1 min, 55°C for 1 min, 72°C for 1 min, 30 cycles) was performed using TaKaRa Taq (Takara Shuzo, Ohtsu) in the Thermal Cycler 480 (Perkin-Elmer, CA, USA).

**[0148]** An about 2 kb fragment obtained was treated with restriction enzymes KpnI and BglII (Takara Shuzo), and was cloned into the KpnI-BglII site of a reporter gene plasmid pGL3-basic (Promega, WI, USA) using the DNA ligation kit ver. II (Takara Shuzo) to transform into competent E. coli JM109 (Nippongene). The transformed E. coli was cultured at 37°C in the LB medium containing 100  $\mu$ g/ml ampicillin, and the plasmid was prepared using the QIAGEN plasmid maxi kit (QIAGEN, Hilden, Germany).

**[0149]** The plasmid HM-2K/GL3 obtained was treated with restriction enzymes KpnI and XhoI, from which a deletion

clone was constructed using the deletion kit (Takara Shuzo) for kilo-sequence to obtain a plasmid HM-493/GL3 containing from the transcription initiation point to - 493 bp upstream. Furthermore, HM-2k/GL3 was treated with restriction enzymes KpnI and AfII, from which a deletion clone was constructed as described above, and HM-151/GL3 and HM-77/GL3 containing from the transcription initiation point to -151 bp or -77 bp upstream were obtained.

**[0150]** For the introduction of the plasmid into the cell, the polyethyleneimine-Transferrinfection Kit (Tf PEI-Kit) (Bender MedSystems, Vienna, Austria) was used, and for the luciferase assay the Dual-Luciferase Reporter Assay System (Promega) was used. The cell line was cultured overnight in RPMI-1640 containing 50  $\mu$ M Deferrioxamine and 10% FBS. In order to form a complex of the plasmid to be introduced with Tf-PEI, a mixture of the reporter gene plasmid at a final concentration of 20  $\mu$ g/ml, 0.4  $\mu$ g/ml of pRL-SV40, and 1  $\mu$ g/ml of Tf-PEI reagent was prepared and was incubated at room temperature for 20 minutes. 5 X 10<sup>5</sup> cells/ml of cells were added at three volumes of the Tf-PEI/plasmid mixture, and was incubated at 37°C for four hours, washed with the medium, and 100  $\mu$ l per well at a concentration of 2 X 10<sup>5</sup> cells/ml was cultured in a 96-well flat-bottomed plate.

**[0151]** IFN- $\alpha$  was added to a final concentration of 0, 10, 100, or 1000 U/ml, which was cultured at 37°C for two days. After the cells were washed in PBS(-), they were dissolved in 20  $\mu$ l of the Passive Lysis Buffer, six  $\mu$ l of which was applied to the C96 White Polysorp Fluoronunc plate (Nunc). Using the Luminoskan (Labsystems), luminescence intensity was measured for Firefly and Renilla at 30  $\mu$ l of the substrate solution and a measurement time of 10 seconds. The measured values were corrected by Firefly/Renilla, and the relative activity was determined with the control (medium) as one.

**[0152]** As a result, the luciferase activity of the reporter was increased in a IFN- $\alpha$  concentration dependent manner for both of the upstream 2 kbp and 493 bp, confirming that the enhanced transcription activity of the promoter region causes the expression induction of the antigen (Figure 3).

**[0153]** Furthermore, the result of an experiment in which the reporter plasmid 151 bp or 77 bp upstream of the transcription initiation point was used, an enhanced luciferase activity by IFN- $\alpha$  stimulation was observed for the reporter plasmid 151 bp upstream. On the other hand, no changes in activity were noted by IFN- $\alpha$  stimulation in the reporter plasmid 77 bp upstream (Figure 4). In the region of 77-151 bp, a sequence having a high homology with GAS element and ISRE was present, and since it is a transcription regulatory factor that is activated in response to IFN- $\alpha$  stimulation, the transcription regulatory factor of the IRF family was shown to be involved in the activity.

#### Example 3. Enhancement of the amount expressed of HM1.24 antigen in myeloma cells by interferon- $\gamma$

**[0154]** According to the method described in Example 1, 1000 U/ml of the natural type interferon- $\gamma$  (R & D System) was used for analysis. As a result, increases in the amount expressed of HM1.24 antigen were observed in both of the myeloma cell line U266 (Figure 5) and the patient's myeloma cells (Figure 6) as for interferon- $\alpha$ .

#### Example 4. Binding of IRF-2 to the HM1.24 promoter region

**[0155]** In order to identify the transcription factor that binds to the HM1.24 promoter region, the Electrophoresis Mobility Shift Assay (EMSA) with the HM1.24 promoter region as the probe was performed as follows to identify IRF-2 as the binding factor.

##### (1) Preparation of nuclear extract

**[0156]** The myeloma cells U266-B1 (ATCC-TIB196) were cultured in the RPMI-1640 medium (GIBCO-BRL) containing 10% FBS (HyClone) at 37°C in a 5% CO<sub>2</sub> incubator. In order to stimulate the cells by interferon- $\alpha$  (IFN- $\alpha$ ) (Pepro Tech EC), IFN- $\alpha$  was added to the medium to a final concentration of 1000 U/ml, and the cells were recovered at 30 minutes, two hours, four hours, and eight hours after the addition. The cells were suspended into cold PBS (-), centrifuged at 1,000 rpm to discard the supernatant, and suspended in a 10 mM Tris, 10 mM NaCl, and 6 mM MgCl<sub>2</sub> solution.

**[0157]** After allowing to stand in ice for five minutes, centrifugation was repeated, and the supernatant was discarded. The cells were suspended into 10 mM Tris, 10 mM NaCl, 6 mM MgCl<sub>2</sub>, 1 mM DTT, 0.4 mM PMSF, 1 mM Na<sub>3</sub>VO<sub>4</sub>. The cells were homogenized on ice using a glass homogenizer, centrifuged at 6000 g for three minutes, and the supernatant was discarded. The cells were suspended into the extraction buffer (20% glycerol, 20 mM HEPES, 420 mM NaCl, 1.5 mM MgCl<sub>2</sub>, 0.2 mM EDTA, 0.2 mM PMSF, 1 mM DTT, 0.1 mM Na<sub>3</sub>VO<sub>4</sub>, 2 mg/ml aprotinin, and 5 mg/ml leupeptin), and then was allowed to stand in ice for 20 minutes. It was centrifuged at 12000 g for 10 minutes, and the supernatant was recovered.

##### (2) Preparation of the labeled probe

**[0158]** As the probe, ISRE2 was constructed that contains sequences (ttcccagaaa (SEQ ID NO: 10) and ggaaact-

gaaact (SEQ ID NO: 11)) having a homology with GAS (IFN- $\gamma$  activation site: the GAS consensus sequence is ttnccnnaa (SEQ ID NO: 8)) and ISRE (IFN- $\alpha$  stimulation response factor: the ISRE consensus sequence is ngaaanngaaact (SEQ ID NO: 9)), at the HM1.24 promoter region. Thus, oligo DNA ISRE-F2 (aatttctgggaaactgaaactgaaacct (SEQ ID NO: 12)) and ISRE-R2 (aattaggtttcagtttcagttccaga (SEQ ID NO: 13)) were mixed and annealed to form a double stranded DNA probe ISRE2.

[0159] Furthermore, oligo DNA adp-1 (catggcatctactctgtagctattgcagagtgcc (SEQ ID NO: 14)) and adp-2 (catggcactctgcaatagtcatacgaagtagatgc (SEQ ID NO: 15)) were mixed and annealed to form an unrelated probe adp. Probes were labeled using the Band Shift Kit (Amersham Pharmacia Biotech) according to the standard protocol. Thus, 50 ng of double stranded DNA constructed as above was subjected to the polymerase reaction of the Klenow fragment in a reaction solution containing [ $\alpha$ - $^{32}$ P]dATP (20  $\mu$ Ci) (Amersham Pharmacia Biotech) at 37°C for one hour. The solution at the end of the reaction was diluted two-fold and then was loaded to the Nick Spin Column (Amersham Pharmacia Biotech), and after centrifugation at 1600 rpm for four minutes, the solution was recovered to prepare a labeled probe.

### (3) Changes with time in the binding factor produced by stimulation with IFN- $\alpha$

[0160] According to the standard protocol of the Band Shift Kit (Amersham Pharmacia Biotech, NJ, USA), the following procedure was performed. To 5  $\mu$ g of the extracts prepared as the time elapse in the above (1), were added 2  $\mu$ l of the 10 X binding buffer (100 mM Tris-HCl, pH 7.5, 500 mM NaCl, 5 mM DTT, 4  $\mu$ l of 50% glycerol, 1  $\mu$ l of 1% NP-40, and 1  $\mu$ l of poly(dI-dC)-poly(dI-dC)), and then 2  $\mu$ l of the  $^{32}$ P labeled ISRE-2 probe prepared in the above (2), to which water was added to a total volume of 20  $\mu$ l, was added and this reaction mixture was incubated at room temperature for 20 minutes to allow for the binding of the possible binding factors that may be present in the above extract and said  $^{32}$ P labeled ISRE-2 probe.

[0161] To 18  $\mu$ l of the reaction mixture was added 2  $\mu$ l of the 10 X stain solution (attached to the kit), which was electrophoresed in 1 X Tris-glycine buffer (25 mM Tris, 190 mM glycine, 1 mM EDTA, pH 8.1) on a 7.5% acrylamide gel, and then, after electrophoresis, the gel was attached to a filter paper to transfer protein to the filter paper. The filter paper dried with a gel drier was exposed to X-ray film to detect signals.

[0162] For comparison, a reaction solution [(NEC-)] to which no extract was added, a reaction solution [0h] to which an extract from the cell culture that was cultured without stimulation by interferon- $\alpha$  was added, a reaction solution {8h (+cold)} in which 50 ng of a nonlabeled ISRE2 probe was added in stead of the labeled probe to the extract of 8 hour-culture, and a reaction solution [8h (+cold unrelated)] in which 50 ng of an unrelated probe adp was added to the extract of 8 hour-culture were prepared, and were processed as described above to detect signals.

[0163] The result is shown in Figure 7. As can be seen from this figure, a substance that binds to a double stranded DNA corresponding to part of the HM1.24 promoter increased with time in the U266-B1 cells cultured under the stimulation by interferon.

### (4) The identification of a transcription factor by reaction with various antibodies

[0164] In a manner similar to that described in the above (1), the myeloma cells U266-B1 (ATCC-TIB196) were cultured for eight hours in the presence of 1000 U/ml of interferon- $\alpha$  to prepare an extract. The following procedure was performed according to the standard protocol of the Band Shift Kit (Amersham Pharmacia Biotech). Thus, 2  $\mu$ g of antibody was added to 5  $\mu$ g of the extract, and incubated at room temperature for 15 minutes to obtain an extract/antibody reaction solution. To 2  $\mu$ l of the 10 X binding buffer attached to the kit, 4  $\mu$ l of 50% glycerol, 1  $\mu$ l of 1% NP-40, and 1  $\mu$ l of Poly(dI-dC)-Poly(dI-dC) were added 2  $\mu$ l of the above extract/antibody reaction solution and 2  $\mu$ l of the labeled probe prepared in the above (2), to which water was added to make the total volume 20  $\mu$ l, and the reaction mixture was incubated at room temperature for 20 minutes.

[0165] The reaction mixture was subjected to electrophoresis as described in the above (3) to detect signals.

[0166] As the above antibody, the following antibodies (all are from Santa Cruz Biotechnology) were used:

- Anti-human STAT1 p84/p91 (E-23): (description) rabbit polyclonal antibody (SC-346X)
- Anti-human STAT2 (C-20): rabbit polyclonal antibody (SC-476X)
- Anti-mouse STAT3 (K-15): rabbit polyclonal antibody (SC-483X)
- Anti-human ISGF-3 $\gamma$  p48 (C-20): rabbit polyclonal antibody (SC-496X)
- Anti-human IRF-1 (C-20): rabbit polyclonal antibody (SC-497X)
- Anti-human IRF-2 (C-19): rabbit polyclonal antibody (SC-498X)
- Anti-mouse ICSAT (M-17): rabbit polyclonal antibody (SC-6059X)

[0167] As a control, a reaction solution that uses an extract of the cells cultured without interferon stimulation [0h];



a reaction solution in which an extract of the cells cultured for eight hours under the stimulation with 1000 U/ml interferon- $\alpha$  was added, and no antibody was added [8h]; a reaction solution in which 50 ng of the nonlabeled ISRE2 probe was added instead of the labeled ISRE2 [8h (+cold)]; and a reaction solution in which 50 ng of the nonlabeled dp probe was added instead of the labeled ISRE2 probe [8h (+unrelated cold)] were prepared and processed as described above.

**[0168]** The result is shown in Figure 8. As can be seen from this figure, it was shown that the component that binds to the labeled ISRE2 probe in the extract from the cells cultured under the stimulation with interferon- $\alpha$  binds only to anti-IRF-2 antibody, and the factor that binds to and thereby activates the HM1.24 promoter is a transcription factor IRF-2.

#### Example 5. Confirmation of the HM1.24 promoter activation with IRF-2

**[0169]** Effect on HM1.24 promoter activity by IRF-2 co-expression was determined by the reporter gene assay using the U266 cells, and it was revealed that IRF-2 actually has the transcription activation activity of the HM1.24 promoter. In the following experiment, a myeloma cell line U266-B1 (ATCC TIB196) was used. The cells were cultured in the RPMI-1640 medium (GIBCO) (referred to hereinafter as the medium) containing 10% FBS (GIBCO BRL) in a 5% CO<sub>2</sub> incubator.

#### (1) Construction of a plasmid containing the HM1.24 promoter region

**[0170]** The gene of the HM1.24 promoter region was obtained by PCR cloning. From human peripheral blood mononuclear cells, genomic DNA was prepared using the DNAzol reagent (GIBCO). With the genomic DNA obtained as the template, using primer HM2k (aaaggtaccagctgtcttctgtctgtcc) (SEQ ID NO: 16) and BST2B (atagtcatacgaagtagatgccatccag) (SEQ ID NO: 17), PCR (94°C for 1 min, 55°C for 1 min, 72°C for 1 min, 30 cycles) was performed using TaKaRa Taq (Takara Shuzo, Ohtsu) in the Thermal Cycler 480 (Perkin-Elmer, CA, USA).

**[0171]** An obtained about 2 kb fragment was treated with restriction enzymes KpnI and BglII (Takara Shuzo), and was cloned into the KpnI-BglII site of a reporter gene plasmid pGL3-basic (Promega, WI, USA) using the DNA ligation kit ver. II (Takara Shuzo) to transform competent E. coli JM109 (Nippongene). The transformed E. coli was cultured at 37°C in the LB medium containing 100  $\mu$ g/ml ampicillin, and a plasmid was prepared using the QIAGEN plasmid maxi kit (QIAGEN, Hilden, Germany).

**[0172]** The plasmid HM-2k/GL3 obtained was treated with restriction enzymes KpnI and XhoI, from which a deletion clone was constructed using the deletion kit (Takara Shuzo) for kilo-sequence to obtain a plasmid HM-491/GL3 containing up to -491 bp upstream of the transcription initiation point. Furthermore, HM-2k/GL3 was treated with restriction enzymes KpnI and AflII, from which a deletion clone was constructed as described above, and HM-151/GL3 containing -151 bp upstream of the transcription initiation point was obtained.

**[0173]** Furthermore, with HM-2k/GL3 as the template, using primer 10S (ttcgggtacctaattaatcctctgcctg) (SEQ ID NO: 18) and GL primer 2 (cttatgttttggcgtcttcca) (SEQ ID NO: 19), PCR (94°C for 1 min, 55°C for 1 min, 72°C for 1 min, 30 cycles) was performed using TaKaRa Taq (Takara Shuzo, Ohtsu) in the Thermal Cycler 480 (Perkin-Elmer, CA, USA). The fragment obtained was treated with restriction enzymes KpnI and BglII (Takara Shuzo), and was cloned into the KpnI-BglII site of a reporter gene plasmid pGL3-basic (Promega, WI, USA) using the ligation high (Toyobo) to transform competent E. coli JM109 (Nippongene).

**[0174]** The transformed E. coli was cultured at 37°C in the LB medium containing 100  $\mu$ g/ml ampicillin, and a plasmid was prepared using the QIAGEN plasmid maxi kit (QIAGEN, Hilden, Germany). A plasmid HM-125/GL3 was thus obtained that contains up to 125 bp upstream of the transcription initiation point. Furthermore, with HM-2k/GL3 as the template, using primer HMP700 (aaaggtaccagagtttacctggtatctgg) (SEQ ID NO: 20) and GL primer 2, PCR was performed in a similar procedure, and by introducing the fragment into the KpnI-BglII site of pGL3-basic, HM-700/GL3 containing up to about 700 bp upstream of the transcription initiation point was obtained.

**[0175]** Furthermore, with HM-2k/GL3 as the template, using primer HMP700 and 11A' (cagaggattaattaggtaccgaaagagaggtgggctttt) (SEQ ID NO: 21), PCR (98°C for 15 seconds, 65°C for 2 seconds, 74°C for 30 seconds, 25 cycles) was performed using the KOD polymerase (Toyobo) in the Thermal Cycler 480 (Perkin-Elmer, CA, USA). The fragment obtained was inserted into the pCR4 Blunt-TOPO vector using the Zero Blunt TOPO PCR cloning kit for sequencing ver. B (Invitrogen). The plasmid obtained was treated with a restriction enzyme KpnI, and an about 550 bp fragment was recovered, which was introduced into the KpnI site of HM-125/GL3 using the ligation high. Thus, dSIRE/GL3 lacking -25 to about -145 upstream of the transcription initiation point was obtained.

#### (2) Construction of IRF-2 expression plasmid

**[0176]** The IRF-2 expression plasmid was constructed as follows. From the U266 cells for which eight hours have

elapsed after stimulation with interferon- $\alpha$  (1000 U/ml), total RNA was extracted using the TRIzol reagent (GIBCO BRL). With RNA obtained by using the First-strand cDNA Synthesis kit (Pharmacia) as the template, using NotI-d(T)<sub>18</sub> as the primer, a reverse transcription reaction was performed at 37°C for one hour. With the cDNA obtained as the template, using IRF2-F2 (ttgtattgtagcgtgaaaaaagc) (SEQ ID NO: 22) and IRF2-R2 (cagctagttcacattatctcgcc) (SEQ ID NO: 23) as primers, PCR (94°C for 45 seconds, 60°C for 45 seconds, 72°C for 60 seconds, 40 cycles) was performed using LA-Taq (Takara Shuzo).

**[0177]** With the PCR reaction as the template, using IRF2-F1 (agagggtaccatgccggtggaaaggatgcg) (SEQ ID NO: 24) and IRF2-R1 (agtcgggtacctaactgctcttgacgcggg) (SEQ ID NO: 25) as primers, PCR (94°C for 45 seconds, 60°C for 45 seconds, 72°C for 60 seconds, 30 cycles) was performed using the KOD polymerase (Toyobo). The fragment obtained was treated with a restriction enzyme KpnI, and then introduced into the KpnI site of an expression plasmid pTracer-CMV (Invitrogen) using the ligation high (Toyobo) to obtain a IRF-2 expression plasmid pIRF-2/Tracer.

### (3) Measurement of the reporter gene activity

**[0178]** For the introduction of the plasmid into the cells, the polyethyleneimine-Transferrinfection Kit (Tf PEI-Kit) (Bender MedSystems, Vienna, Austria) was used, and for the luciferase assay the Dual-Luciferase Reporter Assay System (Promega) was used. The cell line was cultured overnight in RPMI-1640 containing 50  $\mu$ M Deferrioxamine and 10% FBS. In order to form a complex of the plasmid to be introduced with Tf-PEI, a mixture of the reporter gene plasmid at a final concentration of 20  $\mu$ g/ml, 20  $\mu$ g/ml of pIRF-2/Tracer or pTracer-CMV, 0.4  $\mu$ g/ml of pRL-SV40, and 1  $\mu$ g/ml of Tf-PEI reagent was prepared, and was incubated at room temperature for 20 minutes.

**[0179]** 5 X 10<sup>5</sup> cells/ml of cells were added at three times the volume of the Tf-PEI/plasmid mixture, and was incubated at 37°C for four hours, washed with the medium, and 100  $\mu$ l per well at a concentration of 2 X 10<sup>5</sup> cells/ml was cultured in a 96-well flat-bottomed plate. IFN- $\alpha$  was added to a final concentration of 0 or 1000 U/ml, which was cultured at 37°C for two days. After the cells were washed in PBS(-), the cells were dissolved in 20  $\mu$ l of the Passive Lysis Buffer, six  $\mu$ l of which was applied to the C96 White Polysorp Fluoronunc plate (Nunc). Using the Luminoskan (Labsystems), luminescence intensity was measured for Firefly and Renilla at 30  $\mu$ l of the substrate at a measurement time of 10 seconds. The measured values were corrected by Firefly/Renilla to obtain the correct relative activity.

### (4) Result

**[0180]** The HM1.24 promoter reporter plasmid and the IRF-2 expression plasmid were introduced into the U266 cells, and the reporter activity was determined (Figure 9). As a result, the luciferase activity was increased in -700 and -151 containing the ISRE motif sequence that is a IRF-2 binding site by IRF-2 co-expression. On the other hand, no changes in luciferase activity were noted in dISRE/GL3 that lacks the ISRE sequence by IRF-2 co-expression. This result indicated that IRF-2 binds to the ISRE region of the HM1.24 promoter and enhances its transcription activity.

### (5) Confirmation of enhanced expression of HM1.24 antigen by over expression of IRF-2

**[0181]** For changes in the amount expressed of HM1.24 antigen by IRF-2, the IRF-2 expression plasmid (pIRF-2/Tracer) or the control plasmid (pTracer/CMV) is introduced into the U266 cells in the method described above, then cultured for 1-2 days, from which the cells are recovered, and then stained with mouse anti-human HM1.24 antibody as a primary antibody. The cells are washed, and further stained with FITC-labeled anti-mouse IgG antibody as a secondary antibody. After washing the cells, the FITC fluorescence intensity of the cells was measured by a flow cytometer. It is confirmed that in the cells in which the IRF-2 expression plasmid was introduced there are more cells having a high FITC intensity compared to the cells in which the control plasmid was introduced.

### Example 6. Enhancement of the amount expressed of HM1.24 antigen by interferon- $\alpha$ in lymphatic tumor cells and expression induction of HM1.24 antigen in myelocytic leukemia cells

**[0182]** A human myelocytic leukemia cell line HEL (Japanese Cancer Research Resources, Tokyo, Japan) and tumor cells derived from the bone marrow and lymph node of patients with hematopoietic tumors were cultured in the method described in Example 1. Interferon- $\alpha$  used is the same as in Example 1, and the method of stimulation by interferon- $\alpha$  and the method of flow cytometry were performed in a similar manner to that described in Example 1. As the FITC-anti-HM1.24 antibody, FITC-humanized anti-HM1.24 antibody (1 mg/ml) was used (see International Patent Publication WO 98/14580, the versions of the light chain and the heavy chain variable region are RVL<sub>a</sub> and RVH<sub>s</sub>, respectively). Twenty  $\mu$ l of PE-anti-CD33 antibody for the identification of myelocytic leukemia cells, PE-anti CD19 antibody for the identification of B lymphocytic lymphoma, and PE-anti CD4 antibody (all are from Pharmingen, San Diego, CA, USA) for the identification of T lymphocytic lymphoma were added, and double-stained (Figs. 10 to 12).

[0183] As a result, under no stimulation with IFN- $\alpha$ , very little or little expression of HM1.24 antigen on the cell surface of HEL and tumor cells derived from patients with acute myelocytic leukemia was observed, indicating that the expression of HM1.24 antigen under no stimulation with IFN- $\alpha$  is substantially not observed in these tumor cells. On the other hand, in tumor cells derived from patients with acute lymphocytic leukemia, B lymphocytic non-Hodgkin lymphoma, and T lymphocytic non-Hodgkin lymphoma, the expression of HM1.24 antigen on the cell surface was observed even under no stimulation with IFN- $\alpha$ . In contrast, when stimulated with IFN- $\alpha$ , the expression of HM1.24 antigen was induced in HEL and tumor cells derived from patients with acute myelocytic leukemia.

[0184] In these cells that substantially do not express HM1.24 antigen, IFN- $\alpha$  could cause the expression of HM1.24 antigen. Also, in tumor cells derived from patients with acute lymphocytic leukemia, B lymphocytic non-Hodgkin lymphoma, and T lymphocytic non-Hodgkin lymphoma, the amount expressed of HM1.24 antigen increased. In these five types of tumor cells, the amount of HM1.24 antigen expressed by IFN- $\alpha$  stimulation was almost the same. These results indicate that IFN- $\alpha$  exhibits the effect of inducing or enhancing the expression of HM1.24 antigen for hematopoietic tumor cells in general.

#### Example 7. Enhancement of ADCC activity of anti-HM1.24 antibody by interferon $\alpha$

[0185] A human myelocytic leukemia cell line HEL was used as the target cell. To HEL ( $1 \times 10^6$  cells), 0.1 mCi of 51 Cr-sodium chromate (New England Nuclear, Boston, MA, USA) was added and was allowed to stand at 37°C for 1 hour. Then the cells were washed three times in RPMI1640, and  $1 \times 10^4$  cells were dispensed into a roundbottomed 96-well plate (Corning).

[0186] After adding various concentrations of humanized anti-HM1.24 antibody (International Patent Publication WO 98/14580, the versions of the light chain and the heavy chain variable region are RVL $\alpha$  and RVHs, respectively), peripheral blood mononuclear cells ( $5 \times 10^5$ ) from normal healthy subjects were added, and were allowed to stand at 37°C for 4 hours. The radioactivity of the culture supernatant was counted by a gamma counter, and cytotoxicity by ADCC was calculated by the following equation. The maximum value was measured by disrupting the cells by adding 1% NP40 to the target cells. The minimum value was measured by adding the culture liquid RPMI1640 alone (Fig. 13).

$$\text{ADCC activity (\%)} = \frac{(\text{measured value} - \text{minimum value})}{(\text{maximum value} - \text{minimum value})}$$

[0187] As a result, in the absence of stimulation by IFN- $\alpha$ , no concentrations of humanized anti-HM1.24 antibody caused marked ADCC activity. When stimulated by IFN- $\alpha$ , marked ADCC activity due to humanized anti-HM1.24 antibody was noted. Furthermore, ADCC activity was dependent on antibody concentration, which indicated that the expression of HM1.24 antigen on the cell surface induced ADCC activity.

[0188] The result indicates that the enhanced amount expressed of HM1.24 antigen on the surface of tumor cells due to stimulation by IFN- $\alpha$  etc. leads to enhancement in cytotoxicity due to anti-HM1.24 antibody, i.e. enhancement in anti-tumor effect and, therefore, that the combined use of an expression-enhancing agent such as IFN- $\alpha$  of anti-HM1.24 antibody could attain an equivalent effect with the use of a smaller amount of anti-HM1.24 antibody. It also indicates that, even in hematopoietic tumor cells for which a marked effect cannot be expected with anti-HM1.24 antibody alone, the combined use of an expression-enhancing agent such as IFN- $\alpha$  and anti-HM1.24 antibody is likely to attain an anti-tumor effect.

## SEQUENCE LISTING

5 <110> CHUGAI SEIYAKU KABUSHIKI KAISHA

<120> Therapeutic agent for hematopoietic organ tumors

10 <130> H757

<160> 5

15 <210> 1

<211> 1013

<212> DNA

20 <213> Homo sapiens

<223> Nucleotide sequence coding for HM1.24 protein  
antigen

25

<400> 1

gaattcggca cgagggatct gg atg gca tct act tcg tat gac tat tgc 49

30 Met Ala Ser Thr Ser Tyr Asp Tyr Cys

1 5

aga gtg ccc atg gaa gac ggg gat aag cgc tgt aag ctt ctg ctg ggg 97

Arg Val Pro Met Glu Asp Gly Asp Lys Arg Cys Lys Leu Leu Leu Gly

35 10 15 20 25

ata gga att ctg gtg ctc ctg atc atc gtg att ctg ggg gtg ccc ttg 145

Ile Gly Ile Leu Val Leu Leu Ile Ile Val Ile Leu Gly Val Pro Leu

30 35 40

40 att atc ttc acc atc aag gcc aac agc gag gcc tgc cgg gac ggc ctt 193

Ile Ile Phe Thr Ile Lys Ala Asn Ser Glu Ala Cys Arg Asp Gly Leu

45 45 50 55

cgg gca gtg atg gag tgt cgc aat gtc acc cat ctc ctg caa caa gag 241

45 Arg Ala Val Met Glu Cys Arg Asn Val Thr His Leu Leu Gln Gln Glu

60 65 70

50

55

EP 1 364 657 A1

5                    75                    80                    85  
 ctg acc gag gcc cag aag ggc ttt cag gat gtg gag gcc cag gcc gcc 289  
 Leu Thr Glu Ala Gln Lys Gly Phe Gln Asp Val Glu Ala Gln Ala Ala  
 10                    90                    95                    100                    105  
 acc tgc aac cac act gtg atg gcc cta atg gct tcc ctg gat gca gag 337  
 Thr Cys Asn His Thr Val Met Ala Leu Met Ala Ser Leu Asp Ala Glu  
 15                    110                    115                    120  
 aag gcc caa gga caa aag aaa gtg gag gag ctt gag gga gag atc act 385  
 Lys Ala Gln Gly Gln Lys Lys Val Glu Glu Leu Glu Gly Glu Ile Thr  
 20                    125                    130                    135  
 aca tta aac cat aag ctt cag gac gcg tct gca gag gtg gag cga ctg 433  
 Thr Leu Asn His Lys Leu Gln Asp Ala Ser Ala Glu Val Glu Arg Leu  
 25                    140                    145                    150  
 aga aga gaa aac cag gtc tta agc gtg aga atc gcg gac aag aag tac 481  
 Arg Arg Glu Asn Gln Val Leu Ser Val Arg Ile Ala Asp Lys Lys Tyr  
 30                    155                    160                    165  
 tac ccc agc tcc cag gac tcc agc tcc gct gcg gcg ccc cag ctg ctg 529  
 Tyr Pro Ser Ser Gln Asp Ser Ser Ser Ala Ala Ala Pro Gln Leu Leu  
 35                    170                    175                    180  
 att gtg ctg ctg ggc ctc agc gct ctg ctg cag tga gatcccagga 575  
 Ile Val Leu Leu Gly Leu Ser Ala Leu Leu Gln  
 40                    190                    195                    200  
 agctggcaca tcttgggaagg tccgtcctgc tcggcttttc gcttgaacat tcccttgatc 635  
 tcatcagttc tgagcgggtc atggggcaac acggttagcg gggagagcac ggggtagccg 695  
 gagaaggggc tctggagcag gtctggaggg gccatggggc agtcctgggt ctggggacac 755  
 agtcgggttg acccaggggt gtctccctcc agagcctccc tccggacaat gagtcccccc 815  
 45                    205                    210                    215  
 tcttgtctcc caccctgaga ttgggcatgg ggtgcggtgt ggggggcatg tgctgcctgt 875  
 tgttatgggt tttttttgcg ggggggggtg cttttttctg ggtcttttga gctccaaaaa 935  
 aataaacact tcctttgagg gagagcacac cttaaaaaaa aaaaaaaaaa aaaaaaaaaa 995  
 50                    220                    225                    230  
 aaaattcggg cggccgcc 1013  
 <210>            2  
 <211>            180  
 <212>            PRT  
 55                    235                    240  
 <213>            Homo sapiens  
 <223>            Amino acid sequence of HM1.24 protein antigen

&lt;400&gt; 2

5 Met Ala Ser Thr Ser Tyr Asp Tyr Cys Arg Val Pro Met Glu Asp Gly  
 1 5 10 15  
 Asp Lys Arg Cys Lys Leu Leu Leu Gly Ile Gly Ile Leu Val Leu Leu  
 20 25 30  
 10 Ile Ile Val Ile Leu Gly Val Pro Leu Ile Ile Phe Thr Ile Lys Ala  
 35 40 45  
 Asn Ser Glu Ala Cys Arg Asp Gly Leu Arg Ala Val Met Glu Cys Arg  
 50 55 60  
 15 Asn Val Thr His Leu Leu Gln Gln Glu Leu Thr Glu Ala Gln Lys Gly  
 65 70 75 80  
 Phe Gln Asp Val Glu Ala Gln Ala Ala Thr Cys Asn His Thr Val Met  
 85 90 95  
 20 Ala Leu Met Ala Ser Leu Asp Ala Glu Lys Ala Gln Gly Gln Lys Lys  
 100 105 110  
 Val Glu Glu Leu Glu Gly Glu Ile Thr Thr Leu Asn His Lys Leu Gln  
 115 120 125  
 25 Asp Ala Ser Ala Glu Val Glu Arg Leu Arg Arg Glu Asn Gln Val Leu  
 130 135 140  
 Ser Val Arg Ile Ala Asp Lys Lys Tyr Tyr Pro Ser Ser Gln Asp Ser  
 145 150 155 160  
 30 Ser Ser Ala Ala Ala Pro Gln Leu Leu Ile Val Leu Leu Gly Leu Ser  
 165 170 175  
 Ala Leu Leu Gln  
 35 180

&lt;210&gt; 3

&lt;211&gt; 2061

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

45 <223> Nucleotide sequence of promoter region of gene  
 coding for HM1.24 protein antigen

&lt;400&gt; 3

EP 1 364 657 A1

actaaaagtc tctgatatgc agaaataatg gcataagctg tctttctgtc tgtccccctet 60  
 5 ctctctctct gctcgcgctg ccaggcaggg aagggccccc tgtccagtgg acacgtgacc 120  
 cacatgacct tacctatcat tggagatgac tcacactctt taccctgccc cttttgcttt 180  
 gtatccaata aataacagca cagccagaca ttcggggcca ctaccagtct ccgcgcattg 240  
 ctggtagtgg tccccgggc ccagctgtct tttcttttat ctcttcgtct tgtgtcttta 300  
 10 tttctacact ctctcgtcgc cgcacacagg gagagaccca ctgaccctgt ggggctggtc 360  
 cctacagtaa ttttaaaggg aagagcaaca aactttcggg ttgcagggct gggactgttt 420  
 acagctgcaa aatttagaga ggacatcaat ctattattat ccacatttta cagctgggga 480  
 aatcaatgct aagagaggaa attcatttgc ccagagggtgc accaccctgg cctccaatgt 540  
 15 gcaattcatg caattgtgat ttccgacctg gtcccaaaact aaccctaaag ttagcaggcc 600  
 agaacagtgc tgcctcaaata agtcagctta gtcaaataag tcaggcaaag gtcgtgtctt 660  
 tgcacctgga gtcctggcca ggctggtagg tccctcctcc tgggacaagt tcacctcag 720  
 aattttcagc aagatcatct cccacagctt gttaattggg tcttggttct aagtgtttt 780  
 20 tttgtttatt ggtttaagag atgggatccc actctatcac ccaggcttga gtgccgtggc 840  
 acaatcatag ctgcctgcag cctcaaaact ctgggctcga gtgatcctcc tgcctcagcc 900  
 tcccagcctc agcctgggac cacaggcatg taccaccatg cctggctcta agtggtctta 960  
 atggggctct tctgagggat gttggagtca gggcctgggg ggagttcccc aggccttctg 1020  
 25 ggaggcctgg gctctggact tgacctcgcc tactgtctgg ccctgctgaa aagaaaaaaa 1080  
 aacatggaaa tggcagacct aacagaatct gggctgtggg caggatgtgg ctgaagaagc 1140  
 cacaagaaaa acatgcagtc ccctttcagc ggtcatgccc agcagttggg tgccgataat 1200  
 gggcctgatt tctgttagga agcctggct ctcttgcca catggacagt gtctgaggct 1260  
 30 ggccctgtta ttccctttg cagatgaaga aacaggctca gagagtttac ctggtatcct 1320  
 ggagtcccag gagcactttt tctggaagta ggagcttgtt tctgcagggt gccaagacag 1380  
 agaccgacat tgtttgttg ctgggtcggg ctcccagttt tcagctggct ccagtctcac 1440  
 ctgttgetca cacaccctcc atgtctccca tagtcccctc ggtggggaca gaggcactgg 1500  
 35 atgaagccct gctcgtcacc acagagacac ctgaacacaa aaaccagtoc ctggggtcag 1560  
 acccaggccc cgccccaga cccaggccct gccctcactc caccacgcaa ctgtgcaacc 1620  
 tcagtttccc cagggtggaga ccggaccaac aatgatggcc tctgcctctt caggctcatg 1680  
 tacagatgaa tacaggctgg caggccctag gactcagta acacacggca gaggcacagg 1740  
 40 gacttaagat ggagtgtccc aggcagccac agttggctgg caccagttg ggaagggcc 1800  
 aagggctttt aaagcagggt gaaaaaaaaa gccacctcc tttctgggaa actgaaactg 1860  
 aaaacctaata taatcctctg cctgtagggt cctcatgcaa gagctgctgg tcagagcact 1920  
 45 tcttggaact tgctattggt caggacgttt cctatgctaa taaaggggtg gcccgtagaa 1980  
 gattccagca cctccccta actccaggcc agactccttt cagctaaagg ggagatctgg 2040  
 atg gca tct act tcg tat gac 2061  
 Met Ala Ser Thr Ser Tyr Asp

5       <210>       4  
        <211>       29  
        <212>       DNA  
        <213>       Artificial Sequence

10  
        <220>  
        <221>  
        <222>  
   15       <223>       Primer HM2K

       <400>       4  
   20       aaaggtagca gctgtctttc tgtctgtcc                   29

       <210>       5  
        <211>       28  
   25       <212>       DNA  
        <213>       Artificial Sequence

       <220>  
   30       <221>  
        <222>  
        <223>       Primer BST2B

35  
        <400>       5  
        atagtcatac gaagtagatg ccatccag                   28

40       <210>       6  
        <211>       2144  
        <212>       DNA  
        <213>       Homo sapiens

45  
        <223>       Nucleotide sequence coding for IRF-2 protein

50       <400>       6

55



EP 1 364 657 A1

aactgacggg ctttcatttc catttcacac accctagcaa cacttatacc ttgcggaatt 60  
 gtattggtag cgtgaaaaaa gcacactgag agggcaccat gccggtggaa aggatgcgca 120  
 5 tgcccccgtg gctggaggag cagataaact ccaacacgat cccggggctc aagtggctta 180  
 acaaggaaaa gaagattttt cagatcccct ggatgcatgc ggctagacat gggtaggatg 240  
 tggaaaaaga tgcaccactc tttagaaacc gggcaatcca tacaggaaag catcaaccag 300  
 10 gagtagataa acctgatccc aaaacatgga aggcgaattt cagatgcgcc atgaattcct 360  
 tgcctgatat tgaagaagtc aaggataaaa gcataaagaa aggaaataat gccttcaggg 420  
 tctaccgaat gctgccccta tcagaacggc cttctaagaa aggaaagaaa ccaaagacag 480  
 aaaaagaaga caaagttaag cacatcaagc aagaaccagt tgagtcattc ctggggctta 540  
 15 gtaatggagt aagtgatctt tctcctgagt atgcggctct gacttcaact ataaaaaatg 600  
 aagtggatag tacggtgaac atcatagttg taggacagtc ccatctggac agcaacattg 660  
 agaatcaaga gattgtcacc aatccgccag acatttgcca agttgtagag gtgaccactg 720  
 agagcgacga gcagccggctc agcatgagcg agctctaccc tctgcagatc tcccccggtg 780  
 20 ctctctatgc agaaagcgaa acgactgata gtgtgccag cgatgaagag agtgccgagg 840  
 ggccggccaca ctggcggaag aggaatattg aaggcaaaaca gtaoctoagc aacatgggga 900  
 ctcgaggctc ctaccgtctg cccggcatgg cgtctctcgt cacttccaac aaaccggacc 960  
 tccaggtcac catcaaagag gagagcaatc cggtgcctta caacagctcc tggccccctt 1020  
 25 ttcaagacct ccccccttct tctccatga ccccgacac cagcagcagt cggccagacc 1080  
 gggagaccog ggccagcgctc atcaagaaaa catcgatat caccagggcc cgcgtcaaga 1140  
 gctgttaagc ctctgactct ccgcggtggt tgttggggct tcttggcttt gttttgttgt 1200  
 ttgtttgtat tttatttttt tctctctgac acctatttta gacaaatcta agggaaaaag 1260  
 30 ccttgacaat agaacattga ttgctgtgtc caactccagt acctggagct tctctttaac 1320  
 tcaggactcc agcccatggt tagacgtgtg tttctagagc ctgctggatc tcccagggct 1380  
 actcactcaa gttcaaggac caacaagggc agtgagggtg ctgcattgcc tgcggtcaag 1440  
 gccagcaagg tggagtggat gcctcagaac ggacgagata atgtgaacta gctggaattt 1500  
 35 tttattcttg tgaatatgta cataggcagc actagcgaca ttgcagctctg cttctgcacc 1560  
 ttatcttaaa gcacttacag ataggccttc ttgtgatctt gctctatctc acagcacact 1620  
 cagcaccccc ttctctgccc attccccagc ctctcttctt atcccatccc atcccatccc 1680  
 atcccatccc atcccatccc gctcttttcc tacttttctt tccctcaaag cttccattcc 1740  
 40 acatccggag gagaagaagg aaatgaattt ctctacagat gtcccatttt cagactgctt 1800  
 taataaaaaat ccttctaate tgctatgctt gaatgccacg cggtagaaag gaaaaagtat 1860  
 catggaaata ttatgcaaat tcccagattt gaagacaaaa atactctaata tctaaccaga 1920  
 45 gcaagctttt ttatttttta tacaggggaa tttttattc aaggtaaaat tctaaataaa 1980  
 atataattgt tttttatctt ttctacagca aatttataat ttttaagattc cttttcttgt 2040  
 ttatcagcag ttgttattac atccttgttg cacatttttt ttaatttttg taaaggtgaa 2100  
 aaaagctttt atgagctcat ctagcaatca gattttcctg tgga 2144

50

55

<210> 7  
 <211> 349  
 <212> PRT  
 <213> Homo sapiens

<223> Amino acid sequence of IRF-2 protein

<400> 7

Met	Pro	Val	Glu	Arg	Met	Arg	Met	Arg	Pro	Trp	Leu	Glu	Glu	Gln	Ile
1				5					10					15	
Asn	Ser	Asn	Thr	Ile	Pro	Gly	Leu	Lys	Trp	Leu	Asn	Lys	Glu	Lys	Lys
			20					25					30		
Ile	Phe	Gln	Ile	Pro	Trp	Met	His	Ala	Ala	Arg	His	Gly	Trp	Asp	Val
		35					40					45			
Glu	Lys	Asp	Ala	Pro	Leu	Phe	Arg	Asn	Arg	Ala	Ile	His	Thr	Gly	Lys
	50					55					60				
His	Gln	Pro	Gly	Val	Asp	Lys	Pro	Asp	Pro	Lys	Thr	Trp	Lys	Ala	Asn
	65				70				75					80	
Phe	Arg	Cys	Ala	Met	Asn	Ser	Leu	Pro	Asp	Ile	Glu	Glu	Val	Lys	Asp
			85					90						95	
Lys	Ser	Ile	Lys	Lys	Gly	Asn	Asn	Ala	Phe	Arg	Val	Tyr	Arg	Met	Leu
			100					105					110		
Pro	Leu	Ser	Glu	Arg	Pro	Ser	Lys	Lys	Gly	Lys	Lys	Pro	Lys	Thr	Glu
		115					120					125			
Lys	Glu	Asp	Lys	Val	Lys	His	Ile	Lys	Gln	Glu	Pro	Val	Glu	Ser	Ser
		130				135						140			
Leu	Gly	Leu	Ser	Asn	Gly	Val	Ser	Asp	Leu	Ser	Pro	Glu	Tyr	Ala	Val
	145				150				155					160	
Leu	Thr	Ser	Thr	Ile	Lys	Asn	Glu	Val	Asp	Ser	Thr	Val	Asn	Ile	Ile
			165					170					175		
Val	Val	Gly	Gln	Ser	His	Leu	Asp	Ser	Asn	Ile	Glu	Asn	Gln	Glu	Ile
		180					185					190			
Val	Thr	Asn	Pro	Pro	Asp	Ile	Cys	Gln	Val	Val	Glu	Val	Thr	Thr	Glu
		195				200					205				
Ser	Asp	Glu	Gln	Pro	Val	Ser	Met	Ser	Glu	Leu	Tyr	Pro	Leu	Gln	Ile
		210				215					220				

**EP 1 364 657 A1**

```

Ser Pro Val Ser Ser Tyr Ala Glu Ser Glu Thr Thr Asp Ser Val Pro
225                230                235                240
Ser Asp Glu Glu Ser Ala Glu Gly Arg Pro His Trp Arg Lys Arg Asn
                245                250                255
Ile Glu Gly Lys Gln Tyr Leu Ser Asn Met Gly Thr Arg Gly Ser Tyr
                260                265                270
Leu Leu Pro Gly Met Ala Ser Phe Val Thr Ser Asn Lys Pro Asp Leu
                275                280                285
Gln Val Thr Ile Lys Glu Glu Ser Asn Pro Val Pro Tyr Asn Ser Ser
                290                295                300
Trp Pro Pro Phe Gln Asp Leu Pro Leu Ser Ser Ser Met Thr Pro Ala
305                310                315                320
Ser Ser Ser Ser Arg Pro Asp Arg Glu Thr Arg Ala Ser Val Ile Lys
                325                330                335
Lys Thr Ser Asp Ile Thr Gln Ala Arg Val Lys Ser Cys
                340                345

<210>      8
<211>      9
<212>      DNA
<213>      Artificial Sequence

<220>
<221>
<222>
<223>      IFN-gamma activated siRNA (GAS) consensus Sequence

<400>      8
          ttncnnnaa

<210>      9
<211>      13
<212>      DNA
<213>      Artificial Sequence

<220>

```

5 <221>  
 <222>  
 <223> IFN-alpha stimulated response element (ISRE)  
 consensus Sequence

10 <400> 9  
 ngaaanngaa act 13

15 <210> 10  
 <211> 9  
 <212> DNA  
 <213> Artificial Sequence

20 <220>  
 <221>  
 <222>  
 25 <223>

<400> 10  
 ttcccagaa 9

30 <210> 11  
 <211> 13  
 <212> DNA  
 35 <213> Artificial Sequence

40 <220>  
 <221>  
 <222>  
 <223>

45 <400> 11  
 ggaaactgaa act 13

50 <210> 12

55

5 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

10 <220>  
 <221>  
 <222>  
 <223> ISRE-F2 probe

15 <400> 12  
 aatttctggg aaactgaaa tgaaaacct 29

20 <210> 13  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

25 <220>  
 <221>  
 <222>  
 30 <223> ISRE-F2 probe

35 <400> 13  
 aattaggttt tcagtttcag tttcccaga 29

40 <210> 14  
 <211> 37  
 <212> DNA  
 <213> Artificial Sequence

45 <220>  
 <221>  
 <222>  
 <223> adp-1 probe

50

55

	<400>	14	
5	catggcatct acttcgtatg actattgcag agtgcc		37
	<210>	15	
	<211>	36	
10	<212>	DNA	
	<213>	Artificial Sequence	
	<220>		
15	<221>		
	<222>		
	<223>	adp-2 probe	
20	<400>	15	
	catgggcact ctgcaatagt catacgaagt agatgc		36
25	<210>	16	
	<211>	29	
	<212>	DNA	
	<213>	Artificial Sequence	
30	<220>		
	<221>		
	<222>		
35	<223>	Primer HM2k	
	<400>	16	
40	aaaggtacca gctgtcttct tgtctgtcc		29
	<210>	17	
	<211>	28	
45	<212>	DNA	
	<213>	Artificial Sequence	
50	<220>		

55

5           <221>  
           <222>  
           <223>       BST2B

10           <400>       17  
           atagtcatac gaagtagatg ccatccag                   28

15           <210>       18  
           <211>       28  
           <212>       DNA  
           <213>       Artificial Sequence

20           <220>  
           <221>  
           <222>  
           <223>       Primer 10S

25           <400>       18  
           tttcggtaac taattaatcc tctgcctg                   28

30           <210>       19  
           <211>       23  
           <212>       DNA  
           <213>       Artificial Sequence

35           <220>  
           <221>  
           <222>  
           <223>       GL Primer 2

40           <400>       19  
           ctttatgttt ttggcgtctt cca                       23

45           <210>       20  
           <211>       30

50  
 55

	<212>	DNA	
5	<213>	Artificial Sequence	
	<220>		
	<221>		
10	<222>		
	<223>	Primer HMP700	
	<400>	20	
15		aaaggtacca gagtttacct ggtatcctgg	30
	<210>	21	
20	<211>	39	
	<212>	DNA	
	<213>	Artificial Sequence	
25	<220>		
	<221>		
	<222>		
30	<223>	Primer 11A'	
	<400>	21	
		cagaggatta attaggtacc gaaagagagg tgggctttt	35
35	<210>	22	
	<211>	24	
	<212>	DNA	
40	<213>	Artificial Sequence	
	<220>		
	<221>		
45	<222>		
	<223>	Primer IRF2-F2	
50	<400>	22	



ttgtattggt agcgtgaaaa aagc

24

5  
 <210> 23  
 <211> 24  
 <212> DNA  
 10 <213> Artificial Sequence

<220>  
 <221>  
 15 <222>  
 <223> Primer IRF2-R2

20 <400> 23  
 cagctagttc acattatctc gtcc

24

<210> 24  
 25 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

30 <220>  
 <221>  
 <222>  
 35 <223> Primer IRF2-F1

<400> 24  
 agagggtacc atgccggtgg aaaggatgcg

30

40 <210> 25  
 <211> 30  
 <212> DNA  
 45 <213> Artificial Sequence

<220>  
 50 <221>

55

<222>

<223> Primer IRF2-R1

<400> 25

agtcggtacc ttaactgctc ttgacgcggg

30

# Claims

1. An expression-enhancing or expression-inducing agent of a protein (HM1.24 antigen) having the amino acid sequence set forth in SEQ ID NO: 2 in hematopoietic tumor cells, said agent comprising interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein as an active ingredient.

2. The expression-enhancing or expression-inducing agent according to claim 1 wherein said hematopoietic tumor is leukemia, lymphoma, or myeloma.
3. The expression-enhancing or expression-inducing agent according to claim 2 wherein said leukemia is acute myelocytic leukemia, chronic myelocytic leukemia, acute lymphocytic leukemia, or chronic lymphocytic leukemia.
4. The expression-enhancing or expression-inducing agent according to claim 2 wherein said lymphoma is Hodgkin's disease, T lymphocytic non-Hodgkin lymphoma, or B lymphocytic non-Hodgkin lymphoma.
5. The expression-enhancing or expression-inducing agent according to claim 2 wherein said myeloma is multiple myeloma.
6. A therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors comprising, as an active ingredient, (1) interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein, and (2) an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.
7. A therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors comprising, as an active ingredient, an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2, wherein interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein is used in combination.
8. A therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors comprising, as an active ingredient, interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein, wherein an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2 is used in combination.
9. The therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors according to any one of claims 6 to 8 wherein said hematopoietic tumor is leukemia, lymphoma, or myeloma.
10. The therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors according to claim 9 wherein said leukemia is acute myelocytic leukemia, chronic myelocytic leukemia, acute lymphocytic leukemia, or chronic lymphocytic leukemia.
11. The therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors according to claim 9 wherein said lymphoma is Hodgkin's disease, T lymphocytic non-Hodgkin lymphoma, or B lymphocytic non-Hodgkin lymphoma.
12. The therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors according to claim 9 wherein said myeloma is multiple myeloma.
13. The therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors according to any one of claims 6 to 8 wherein said antibody is an antibody having cytotoxicity.
14. The therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors according to claim 13 wherein said cytotoxicity is ADCC activity.
15. The therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors according to any one of claims 6 to 14 wherein said antibody is a monoclonal antibody.
16. The therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors according to claim 15 wherein said antibody is a chimeric antibody, a humanized antibody, or a human antibody.
17. The therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors according to claim 15 wherein said antibody is anti-HM1.24 antibody produced by a hybridoma having the Deposit No. FERM BP-5233.
18. The therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors according to claim 16 wherein said chimeric antibody or humanized antibody is a chimeric antibody or humanized antibody of anti-HM1.24 antibody produced by a hybridoma having the Deposit No. FERM BP-5233.

19. A kit for the treatment of a patient having a hematopoietic tumor, said kit comprising:

- (1) an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2;  
and
- (2) an instruction manual instructing the administration of the above antibody to the patient in combination with an expression-enhancing agent of a protein having the amino acid sequence set forth in SEQ ID NO: 2.

20. A kit for the treatment of a patient having a hematopoietic tumor, said kit comprising:

- (1) an expression-enhancing agent of a protein having the amino acid sequence set forth in SEQ ID NO: 2; and
- (2) an instruction manual instructing the administration of the above antibody to the patient in combination with an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.

21. A kit for the treatment of a patient having a hematopoietic tumor, said kit comprising:

- (1) an expression-enhancing agent of a protein having the amino acid sequence set forth in SEQ ID NO: 2; and
- (2) an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2;  
and
- (3) an instruction manual instructing the combined administration of the above agent and the above antibody to the patient.

22. A method of screening an expression-enhancing agent of HM1.24 antigen, comprising the steps of:

- (1) preparing cells having a reporter gene that has the region of the HM1.24 gene promoter;
- (2) contacting said cells with a test substance; and
- (3) detecting the expression of the reporter gene.

23. The expression-enhancing agent of HM1.24 antigen selected by the method according to claim 22.

24. A pharmaceutical composition for the treatment of hematopoietic tumors wherein said composition comprises the expression-enhancing agent according to claim 23 and said composition is used in combination with an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.

25. A method of screening an expression-enhancing agent of the IRF-2 protein comprising the steps of:

- (1) contacting cells with a test substance; and
- (2) determining the amount expressed of the IL-2 protein in said cells.

26. The expression-enhancing agent for the IRF-2 protein, selected by the method according to claim 25.

27. A pharmaceutical composition for the treatment of hematopoietic tumors wherein said composition comprises the expression-enhancing agent according to claim 26 and composition is used in combination with an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.

28. The use of interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein for the production of the expression-enhancing or expression-inducing agent for a protein (HM1.24 antigen) having the amino acid sequence set forth in SEQ ID NO: 2 in hematopoietic tumor cells.

29. The use according to claim 28 wherein said hematopoietic tumor is leukemia, lymphoma, or myeloma.

30. The use according to claim 29 wherein said leukemia is acute myelocytic leukemia, chronic myelocytic leukemia, acute lymphocytic leukemia, or chronic lymphocytic leukemia.

31. The use according to claim 29 wherein said lymphoma is Hodgkin's disease, T lymphocytic non-Hodgkin lymphoma, or B lymphocytic non-Hodgkin lymphoma.

32. The use according to claim 29 wherein said myeloma is multiple myeloma.

33. The use of (1) interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein, and (2) an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2, for the production of a therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors.
- 5 34. The use of an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2, for the production of a therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors, which is used in combination with interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein.
- 10 35. The use of interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein for the production of a therapeutic agent or a pharmaceutical composition for the treatment of hematopoietic tumors, which is used in combination with an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.
- 15 36. The use according to any one of claims 33 to 35 wherein said hematopoietic tumor is leukemia, lymphoma, or myeloma.
37. The use according to claim 36 wherein said leukemia is acute myelocytic leukemia, chronic myelocytic leukemia, acute lymphocytic leukemia, or chronic lymphocytic leukemia.
- 20 38. The use according to claim 36 wherein said lymphoma is Hodgkin's disease, T lymphocytic non-Hodgkin lymphoma, or B lymphocytic non-Hodgkin lymphoma.
39. The use according to claim 36 wherein said myeloma is multiple myeloma.
- 25 40. The use according to any one of claims 33 to 35 wherein said antibody is an antibody having cytotoxicity.
41. The use according to claim 40 wherein said cytotoxicity is ADCC activity.
42. The use according to any one of claims 33 to 41 wherein said antibody is a monoclonal antibody.
- 30 43. The use according to claim 42 wherein said antibody is a chimeric antibody, a humanized antibody, or a human antibody.
44. The use according to claim 45 wherein said antibody is anti-HM1.24 antibody produced by a hybridoma having the Deposit No. FERM BP-5233.
- 35 45. The use according to claim 43 wherein said chimeric antibody or humanized antibody is a chimeric antibody or humanized antibody of anti-HM1.24 antibody produced by a hybridoma having the Deposit No. FERM BP-5233.
- 40 46. The use of the expression-enhancing agent according to claim 23, for the production of a pharmaceutical composition for the treatment of hematopoietic tumors, which is used in combination with an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.
- 45 47. The use of the expression-enhancing agent according to claim 26, for the production of a pharmaceutical composition for the treatment of hematopoietic tumors, which is used in combination with an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.
48. A method of enhancing or inducing the expression of a protein (HM1.24 antigen) having the amino acid sequence set forth in SEQ ID NO: 2 which method comprises administering interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein.
- 50 49. The method according to claim 48 wherein said hematopoietic tumor is leukemia, lymphoma, or myeloma.
50. The method according to claim 49 wherein said leukemia is acute myelocytic leukemia, chronic myelocytic leukemia, acute lymphocytic leukemia, or chronic lymphocytic leukemia.
- 55 51. The method according to claim 49 wherein said lymphoma is Hodgkin's disease, T lymphocytic non-Hodgkin lymphoma, or B lymphocytic non-Hodgkin lymphoma.
52. The method according to claim 49 wherein said myeloma is multiple myeloma.

**53.** A method of treating hematopoietic tumors which comprises administering (1) interferon  $\alpha$ , interferon  $\gamma$ , or the IRF-2 protein, and (2) an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.

**54.** The method according to claim 53 wherein said hematopoietic tumor is leukemia, lymphoma, or myeloma.

**55.** The method according to claim 53 wherein said leukemia is acute myelocytic leukemia, chronic myelocytic leukemia, acute lymphocytic leukemia, or chronic lymphocytic leukemia.

**56.** The method according to claim 53 wherein said lymphoma is Hodgkin's disease, T lymphocytic non-Hodgkin lymphoma, or B lymphocytic non-Hodgkin lymphoma.

**57.** The method according to claim 53 wherein said myeloma is multiple myeloma.

**58.** The method according to claim 53 wherein said antibody is an antibody having cytotoxicity.

**59.** The method according to claim 53 wherein said cytotoxicity is ADCC activity.

**60.** The use according to any one of claims 53 to 59 wherein said antibody is a monoclonal antibody.

**61.** The method according to claim 60 wherein said antibody is a chimeric antibody, a humanized antibody, or a human antibody.

**62.** The method according to claim 60 wherein said antibody is anti-HM1.24 antibody produced by a hybridoma having the Deposit No. FERM BP-5233.

**63.** The method according to claim 61 wherein said chimeric antibody or humanized antibody is a chimeric antibody or humanized antibody of anti-HM1.24 antibody produced by a hybridoma having the Deposit No. FERM BP-5233.

**64.** The method of treating hematopoietic tumors comprising the combined use of the expression-enhancing agent according to claim 23 and an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.

**65.** The method of treating hematopoietic tumors comprising the combined use of the expression-enhancing agent according to claim 26 and an antibody that specifically binds to a protein having the amino acid sequence set forth in SEQ ID NO: 2.

Fig.1





Fig.2







Fig.3




Fig.4




Fig.5





Fig.6





Fig.7




Fig. 8




Fig.9




Fig.10






Fig.11




Fig.12





Fig.13



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/00989

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> Int.Cl <sup>7</sup> A61K38/21, 38/17, 39/395, 45/00, A61P35/00, 35/02, 19/00, C12Q1/02, 1/66, 1/68, C12N15/12, C07K14/705, 16/28  According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) Int.Cl <sup>7</sup> A61K38/21, 38/17, 39/395, 45/00, A61P35/00, 35/02, 19/00, C12Q1/02, 1/66, 1/68, C12N15/12, C07K14/705, 16/28  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAPLUS (STN), MEDLINE (STN), EMBASE (STN)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	OZAKI S. et al., Interferon-alpha and -gamma enhance the HM1.24 expression of myeloma cells through the STAT-signaling pathway, Blood, 1999, Vol.94, No.10, Suppl. 1, Part 1, page 549a	1-24, 27, 28-46, 47
X	WO 94/06818 A1 (Tadatsugu TANIGUCHI) 31 March, 1994 (31.03.94), & EP 662976 A1 & US 5652095 A & US 5807836 A & JP 8-501691 A	25, 26
X	KOENING Merediz S. A. et al., Cloning of an interferon regulatory factor 2 isoform with different regulatory ability, Nucleic Acids. Res., 2000, Nov. 1, Vol.28, No.21, pages 4219 to 4224	25, 26
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 02 May, 2002 (02.05.02)		Date of mailing of the international search report 21 May, 2002 (21.05.02)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/00989

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99/18997 A1 (Chugai Seiyaku Kabushiki Kaisha), 22 April, 1999 (22.04.99), Especially, claims 15, 19 & EP 1023906 A1	1-24, 27, 28-46, 47
A	OZAKI S. et al., Humanized anti-HM1.24 antibody Mediates myeloma cell cytotoxicity that is enhanced by cytokine stimulation of effector cells, Blood, 1999, Vol.93, No.11, pages 3922 to 3930	1-24, 27, 28-46, 47
A	EP 997152 A1 (Chugai Seiyaku Kabushiki Kaisha), 03 May, 2000 (03.05.00), & WO 98/35698 A1 & JP 10-286088 A	1-24, 27, 28-46, 47
P, X	WO 01/13940 A1 (Chugai Seiyaku Kabushiki Kaisha), 01 March, 2001 (01.03.01), (Family: none)	1-24, 27, 28-46, 47
P, X	WO 01/97844 A1 (Idex Pharmaceuticals Corp.), 27 December, 2001 (27.12.01), (Family: none)	1-24, 27, 28-46, 47

Form PCT/ISA/210 (continuation of second sheet) (July 1998)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/00989

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

- 1.
- ☒
- Claims Nos.: 48-65

because they relate to subject matter not required to be searched by this Authority, namely:

The inventions as set forth in claims 48 to 65 pertain to methods for treatment of the human body by therapy and thus relates to a subject matter which this International Searching Authority is not required, under the provisions of Article 17(2)(a)(i) of the PCT and Rule 39.1(iv) of the Regulations

- 2.
- ☐
- Claims Nos.:

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

- 3.
- ☐
- Claims Nos.:

because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6A(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:  
(See extra sheet.)

- 1.
- ☒
- As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

- 2.
- ☐
- As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

- 3.
- ☐
- As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

- 4.
- ☐
- No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.☒ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1998)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/00989

Continuation of Box No.I-1 of continuation of first sheet(1)

under the PCT, to search.

Continuation of Box No.II of continuation of first sheet(1)

The inventions as set forth in claims 1 to 5 and 28 to 32 relate to Hm1.24 antigen expression inducers or potentiators in tumor cells in hematopoietic organs which contain as the active ingredient interferon  $\alpha$ , interferon  $\gamma$  or IRF-2 protein.

The inventions as set forth in claims 6 to 21 and 33 to 45 relate to remedies or medicinal compositions for tumor in hematopoietic organs comprising the above-described interferon  $\alpha$ , interferon  $\gamma$  or IRF-2 protein per se combined with an antibody against Hm1.24.

The invention as set forth in claim 22 relates to a method of screening an Hm1.24 antigen expression potentiator, and the substances selected by the invention as set forth in claim 22 are used in the inventions as set forth in claims 23, 24 and 46.

In contrast thereto, the invention as set forth in claim 25 relates to a method of screening an IRF-2 protein expression potentiator but the Hm1.24 antigen is not essentially required in achieving this invention.

The substances selected by the invention as set forth in claim 25 are used in the inventions as set forth in claims 26, 27 and 47.

Such being the case, the inventions as set forth in claims 1 to 24 and 28 to 46 and the inventions as set forth in claims 25 to 27 and 47 are not considered as relating to a group of inventions so linked as to form a single general inventive concept.